

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

Re PCT/PTO JUN 2005

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/00, A61K 47/48	A2	(11) International Publication Number: WO 98/02454 (43) International Publication Date: 22 January 1998 (22.01.98)
<p>(21) International Application Number: PCT/EP97/03715</p> <p>(22) International Filing Date: 8 July 1997 (08.07.97)</p> <p>(30) Priority Data: 9614871.3 15 July 1996 (15.07.96) GB</p> <p>(71) Applicant (for all designated States except US): ADPROTECH PLC [GB/GB]; Unit 3, Number 2 Orchard Road, Royston, Herts. SG8 5HD (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SMITH, Richard, Anthony, Godwin [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). DODD, Ian [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). MOSSAKOWSKA, Danuta, Ewa, Irena [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).</p> <p>(74) Agent: DAVIES, Jonathan, Mark; Reddie & Grose, 16 Theobalds Road, London WC1X 8PL (GB).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>	
(54) Title: CONJUGATES OF SOLUBLE PEPTIDIC COMPOUNDS WITH MEMBRANE-BINDING AGENTS		
<p>(57) Abstract</p> <p>Soluble derivatives of soluble polypeptides incorporating membrane binding elements, their use in therapy and methods and intermediates including peptide membrane binding elements.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CONJUGATES OF SOLUBLE PEPTIDIC COMPOUNDS WITH MEMBRANE-BINDING AGENTS

This invention relates to polypeptide derivatives, their use in therapy and methods and intermediates for their production.

Essentially all protein drugs are administered as solutions and function *in vivo* in the solution phase. In biochemistry and pharmacology, however, a large number of control and mediator proteins are associated with or function within or on the plasma membranes of cells. Except for soluble, truncated versions of one class of these molecules, no membrane-associated proteins have been developed as therapeutic agents. There are two main reasons for this situation. Firstly, overexpression of proteins that are retained in the membranes of the producer cells is limited by the the low capacity of membranes for proteins and often by the toxic effects of retention when expression is intrinsically efficient. Secondly, extraction of these proteins from membranes requires detergents or organic solvents, often results in inactivation of the protein, leads to difficulties in achieving the high purity needed for drug use and usually gives a product which is hard to formulate for intravenous administration. In addition, retention of very hydrophobic membrane anchoring elements may cause proteins to associate strongly with lipid-binding proteins in blood when administered intravenously thus preventing access to cell membranes.

Soluble, truncated versions of membrane-associated proteins overcome the production difficulties associated with full length proteins. However such truncated molecules lack the membrane binding capability and specificity of the full length proteins which properties may be advantageous or even essential to the desired therapeutic activity.

The main classes of interaction of proteins with membranes can be summarised as follows:

1. Direct and specific interactions with phospholipid head groups or with other hydrophilic regions of complex lipids or indirectly with proteins already inserted in the membrane. The latter may include all the types of intrinsic membrane protein noted below and such interactions are usually with extracellular domains or sequence loops of the membrane proteins;
2. Through anchoring by a single hydrophobic transmembrane helical region near the terminus of the protein. These regions commonly present a hydrophobic face around the entire circumference of the helix cylinder and transfer of this structure to the hydrophilic environment of bulk water is energetically unfavourable.
3. Further anchoring is often provided by a short sequence of generally cationic aminoacids at the cytoplasmic side of the membrane, C-terminal to the transmembrane helix;

4. Through the use of multiple (normally 2-12 and commonly 4,7 and 10) transmembrane regions which are usually predicted to be helical or near-helical. Although these regions are normally hydrophobic overall, they frequently show some amphipathic behaviour - an outer hydrophobic face and an inner more hydrophilic one being identifiable within a helix bundle located in the lipid bilayer;
5. Through postranslationally linked phosphatidyl inositol moieties (GPI-anchors). These are generated by a specific biosynthetic pathway which recognises and removes a specific stretch of C-terminal aminoacids and creates a membrane-associating diacyl glycerol unit linked *via* a hydrophilic carbohydrate spacer to the polypeptide;
- 10 6. In a related process, single fatty acid groups such as myristoyl, palmitoyl or prenyl may be attached postranslationally to one or more sites in a protein (usually at N- or C-termini). Again, amino acids (such as the C-terminal CAAX box in *Ras* proteins) may be removed.

Artificial membranes are considered to be lipid complexes that mimic the basic
15 properties of the cell membrane, i.e., a lipid vacuole with an aqueous interior and a surface chemistry that resembles the cell membrane. The artificial membrane typically contains phospholipids or mimics thereof and may be unilemmellar or bilemmellar and the outer surface will contain charged groups similar to the choline groups of the most abundant phospholipid. The prototype artificial membrane is known as a liposome and
20 the technologies for the construction of liposomes including the incorporation of therapeutically useful agents into them is well known to those in the art. Liposomes have been evaluated in a number of disease states and liposomes containing the anti-fungal Amphotericin are commercially available. In addition, proteoliposomes have been described. For example, the use of immunoliposomes encapsulating amphotericin
25 B has been reported to be of benefit in the treatment of experimental fungal infections in animal models (e.g. Hospenthal, D. et al (1989) *J. Med. Microbiol.* **30** 193-197; Dromer, F. et al (1990) *Antimicrob. Agents Chemother.* **34** 2055-2060).

Mimics of natural or artificial membranes are often related in structure and will mimic one or more properties of the membrane. One such example is the provision of
30 an artificial surface having pendant groups which mimic the phospholipid zwitterionic groups which are found on the outside of cell surfaces. For example WO92/06719 (Biocompatibles Limited) discloses natural and synthetic phospholipids which may be coated on an artificial surface, e.g. a device which, in use, will come into contact with protein-containing or biological fluids, to provide improved biocompatibility and
35 haemocompatibility and WO 94/16749 discloses additional zwitterionic groups that may be used to improve biocompatibility in a similar way.

The present invention provides a soluble derivative of a soluble polypeptide, said derivative comprising two or more heterologous membrane binding elements with low membrane affinity covalently associated with the polypeptide which elements are capable of interacting, independently and with thermodynamic additivity, with components of cellular or artificial membranes exposed to extracellular fluids.

By 'heterologous' is meant that the elements are not found in the native full length protein from which a soluble protein may be derived.

By 'soluble polypeptide' is meant a truncated derivative of a full length protein which lacks its natural membrane binding capability, and/or a polypeptide which has a solubility level in aqueous media of $> 100\mu\text{g/ml}$.

By 'membrane binding element with low membrane affinity' is meant that the element has only moderate affinity for membranes, that is a dissociation constant greater than $0.1\mu\text{M}$, preferably $1\mu\text{M}$ - 1mM . The elements preferably have a size $<5\text{kDa}$.

The derivative should incorporate sufficient elements with low affinities for membrane components to result in a derivative with a high (preferably $0.01 - 10\text{nM}$ dissociation constant) affinity for specific membranes. The elements combine so as to create an overall high affinity for the particular target membrane but the combination lacks such high affinity for other proteins for which single elements may be (low-affinity) ligands.

The elements should be chosen so as to retain useful solubility in pharmaceutical formulation media, preferably $>100\mu\text{g/ml}$. Preferably at least one element is hydrophilic.

The invention thus promotes localisation of a therapeutic protein at cellular membranes and thereby provides one or more of several biologically significant effects with potential therapeutic advantages including:

Potency: If the protein is a receptor and an agonist or antagonist activity is localised on the same surface as the receptor itself, an increase in effective concentration may result from the reduction in the diffusional degrees of freedom.

Pharmacokinetics and dosing frequency: Interaction of a derivatised protein with long-lived cell types or serum proteins would be expected to prolong the plasma residence time of the protein and produce a depot effect through deposition on cell surfaces.

Specificity: Many clinically important pathological processes are associated with specific cell types and tissues (for example the vascular endothelium and the recruitment thereto of neutrophils bearing the sialyl Lewis^x antigen to ELAM-1, see below). Hence targeting the modified protein to regions of membrane containing

pathology-associated membrane markers may improve the therapeutic ratio of the protein targeted.

The derivatives of the invention may be used in association with artificial membranes or mimics thereof to allow delivery of the therapeutic protein to sites where it will provide therapeutic benefit. For example, polypeptides associated with liposomes formed by contacting liposomes with a derivative of the invention may be more stable than the free polypeptide. The liposome may incorporate a therapeutic agent, for example an antiinflammatory or cytotoxic agent. The polypeptide derivative of the invention may thus be used to target the therapeutic agent. When the polypeptide is itself a therapeutic agent, the liposome incorporated therapeutic agent may be used to enhance further the efficacy or tolerability of the therapy.

Association of derivatives of the invention with mimics of cell membranes may be used to concentrate the therapeutic protein at sites where therapeutically useful concentrations of underivatised protein might be difficult to achieve. For example, indwelling medical devices coated with mimics of the phospholipid zwitterionic groups which are found on the outside of cell surfaces, such as those disclosed in WO92/06719 and WO 94/16749, may be additionally treated with derivatives of the invention. For example complement inhibitors derivatised in accordance with the invention could be incorporated into the outer surface of indwelling catheters or hip replacements or heart valves in order to minimise development of inflammation associated with these operations.

It will be appreciated that all associations of heterologous amino acid sequences with a polypeptide which is a soluble derivative of a human protein will need to be assessed for potential immunogenicity, particularly where the amino acid sequence is not derived from a human protein. The problem can be minimised by using sequences as close as possible to known human ones and through computation of secondary structure and antigenicity indices.

Examples of protein therapeutic agents which may be modified according to the invention include but are not restricted to the following:

Base Protein	Cell Target	Therapeutic Application
IL-4 Y124D mutein	B-cells	Anti-allergy (IL-4 antagonist)
Plasminogen activators e.g. Prourokinase, streptokinase, tissue-type plasminogen activator, reteplase	Erythrocytes, vascular endothelium	Prevention of venous thrombosis
Leptin	Choroid plexus, Hypothalamus	Weight loss (agonist)
Complement inhibitors*	Vascular endothelium, Myocytes, Erythrocytes, Lymphocytes	Ischaemic injury, transplantation, inflammation
scFv antibody against cytokines (IL-1, IL-, IL-5, IL-6)	Eosinophils	Asthma, allergic disease
Protein C	Vascular endothelium	Prevention of venous thrombosis
Antibodies against CD4, B7/CD28, CD3/TCR, CD11b(CR3)	Lymphocytes	Immunosuppression
Interferon- β and derivatives	Lymphocytes	Immunomodulation, multiple sclerosis

*Complement regulatory proteins e.g.: CR1 (CD35); DAF (CD55); MCP (CD46); CD59; Factor H; and C4 binding protein; and hybrids or muteins thereof such as CR1-CD59 (S.G.El Feki and D.T.Fearon *Molecular Immunology* 33 (supp 1). p 57, 1996), MCP-DAF (P.J.Higgins *et al*, *J.Immunology*. 158, 2872-2881,1997) and soluble CR1 polypeptide fragments.

The derivative preferably comprises two to eight, more preferably two to four membrane binding elements.

Membrane binding elements are preferably selected from: fatty acid derivatives such as fatty acyl groups; basic amino acid sequences; ligands of known integral membrane proteins; sequences derived from the complementarity-determining region of monoclonal antibodies raised against epitopes of membrane proteins; membrane binding sequences identified through screening of random chemical or peptide libraries.

The selection of suitable combination of membrane binding elements will be guided by the nature of the target cell membrane or components thereof.

Suitable fatty acid derivatives include myristoyl (12 methylene units) which is insufficiently large or hydrophobic to permit high affinity binding to membranes. Studies with myristoylated peptides (eg R.M.Peitzsch & S.McLaughlin, *Biochemistry*, 32, 10436-10443, 1993)) have shown that they have effective dissociation constants with
 5 model lipid systems of $\sim 10^{-4}$ M and around 10 of the 12 methylene groups are buried in the lipid bilayer. Thus, aliphatic acyl groups with about 8 to 18 methylene units, preferably 10-14, are suitable membrane binding elements. Other examples of suitable fatty acid derivatives include long-chain (8-18, preferably 10-14 methylene) aliphatic amines and thiols, steroid and farnesyl derivatives.

10 Membrane binding has been found to be associated with limited (single-site) modification with fatty acyl groups when combined with a cluster of basic aminoacids in the protein sequence which may interact with acidic phospholipid head groups and provide the additional energy to target membrane binding. This combination of effects has been termed the 'myristoyl-electrostatic switch' (S.McLaughlin and A.Aderem, *TIBS*,
 15 20,272-276, 1994; J.F.Hancock *et al*, *Cell*, 63, 133-139,1990). Thus, a further example of suitable membrane binding elements are basic aminoacid sequences such as those found in proteins such as *Ras* and MARCKS (myristoylated alanine-rich C-kinase substrate, P.J. Blackshear, *J. Biol. Chem.*, 268, 1501-1504, 1993) which mediate the electrostatic 'switch' through reversible phosphorylation of serine residues within the
 20 sequence and a concomitant neutralisation of the net positive charge. Such sequences include but are not restricted to consecutive sequences of Lysine and Arginine such as (Lys)_n where n is from 3 to 10, preferably 4 to 7.

Suitable examples of amino acid sequences comprising basic amino acids include:

- i) DGPKKKKKKSPSKSSG
- 25 ii) GSSKSPSKKKKKKPGD
- iii) SPSNETPKKKKKRFSFKKSG
- iv) DGPKKKKKKSPSKSSK
- v) SKDGKKKKKKSKTK
- (N-terminus on left)

30 Sequences i) to v) are examples of electrostatic switch sequences.

Examples of amino acid sequences derived from ligands of known integral membrane proteins include RGD-containing peptides such as GRGDSP which are ligands for the $\alpha_v\beta_3$ integrin of human platelet membranes. Another example is DGPSEILRGDFSS derived from human fibrinogen alpha chain, which binds to the
 35 GpIIb/IIIa membrane protein in platelets.

Further examples of such sequences include those known to be involved in interactions between membrane proteins such as receptors and the major

histocompatibility complex. An example of such a membrane protein ligand is the sequence GNEQSFRVDLRTLRYA which has been shown to bind to the major histocompatibility complex class 1 protein (MHC-1) with moderate affinity (L.Olsson *et al*, Proc. Natl .Acad.Sci.USA. 91, 9086-909, 1994).

5 Yet further examples of such sequences employ a membrane insertive address specific for T-cells. Such sequence is derived from the known interaction of the transmembrane helix of the T-cell antigen receptor with CD3 (Nature Medicine 3, 84-88,1997). Examples are peptides containing the sequence GFRILLKV such as:

SAAPSSGFRILLKV


10 AAPSVIGFRILLKVAG

An example of a ligand for an integral membrane protein is the carbohydrate ligand Sialyl Lewis^x which has been identified as a ligand for the integral membrane protein ELAM-1 (M.L.Phillips *et al*, Science, 250, 1130-1132, 1990 & G.Walz *et al*, *Ibid*, 250, 1132-1135,1990).

15 Sequences derived from the complementarity-determining regions of monoclonal antibodies raised against epitopes within membrane proteins (see, for example, J.W.Smith *et al*, J.Biol.Chem. 270, 30486-30490, 1995) are also suitable membrane binding elements, as are binding sequences from random chemical libraries such as those generated in a phage display format and selected by biopanning operations *in vitro* (G.F.Smith and J.K.Scott, Methods in Enzymology, 217H, 228-257,1993) or *in vivo* (R.Pasqualini & E.Ruoslahti, Nature, 380, 364-366, 1996).

20 Optionally, conditional dissociation from the membrane may be incorporated into derivatives of the invention using mechanisms such as pH sensitivity (electrostatic switches), regulation through metal ion binding (using endogenous Ca²⁺, Zn²⁺ and incorporation of ion binding sites in membrane binding elements) and protease cleavage (e.g plasminolysis of lysine-rich membrane binding sequences to release and activate prourokinase)

Preferred derivatives of this invention have the following structure:



30

in which:

P is the soluble polypeptide,

each **L** is independently a flexible linker group,

each **W** is independently a peptidic membrane binding element,

35

n is an integer of 1 or more and

X is a peptidic or non-peptidic membrane-binding entity which may be covalently linked to any **W**.

Peptidic membrane binding elements are preferably 8 to 20 amino acids long and elements W are preferably located sequentially either at the N or C terminus of the soluble polypeptide. The amino acid sequences are linked to one another and to the soluble peptide by linker groups which are preferably selected from hydrophilic and/or flexible aminoacid sequences of 4 to 20 aminoacids; linear hydrophilic synthetic polymers; and chemical bridging groups.

Peptide linkages may be made chemically or biosynthetically by expression of appropriate coding DNA sequences. Non peptide linkages may be made chemically or enzymatically by post-translational modification.

The polypeptide portion of the derivatives of the invention may be prepared by expression in suitable hosts of modified genes encoding the soluble polypeptide of interest plus one or more peptidic membrane binding elements and optional residues such as cysteine to introduce linking groups to facilitate post translational derivatisation with additional membrane binding elements.

In a further aspect, therefore, the invention provides a process for preparing a derivative according to the invention which process comprises expressing DNA encoding the polypeptide portion of said derivative in a recombinant host cell and recovering the product and thereafter post translationally modifying the polypeptide to chemically introduce membrane binding elements.

In particular, the recombinant aspect of the process may comprise the steps of:

- i) preparing a replicable expression vector capable, in a host cell, of expressing a DNA polymer comprising a nucleotide sequence that encodes said polypeptide portion;
- ii) transforming a host cell with said vector;
- iii) culturing said transformed host cell under conditions permitting expression of said DNA polymer to produce said polypeptide; and
- iv) recovering said polypeptide.

Where the polypeptide portion is novel, the DNA polymer comprising a nucleotide sequence that encodes the polypeptide portion as well as the polypeptide portion itself and S-derivatives thereof, also form part of the invention. In particular the invention provides a polypeptide portion of a derivative of the invention comprising the soluble peptide linked by a peptide bond to one peptidic membrane binding element and/or including a C-terminal cysteine, and DNA polymers encoding the polypeptide portion.

The recombinant process of the invention may be performed by conventional recombinant techniques such as described in Sambrook *et al.*, Molecular Cloning : A

laboratory manual 2nd Edition. Cold Spring Harbor Laboratory Press (1989) and DNA Cloning vols I, II and III (D. M. Glover ed., IRL Press Ltd).

The invention also provides a process for preparing the DNA polymer by the condensation of appropriate mono-, di- or oligomeric nucleotide units.

5 The preparation may be carried out chemically, enzymatically, or by a combination of the two methods, *in vitro* or *in vivo* as appropriate. Thus, the DNA polymer may be prepared by the enzymatic ligation of appropriate DNA fragments, by conventional methods such as those described by D. M. Roberts *et al.*, in *Biochemistry* 1985, 24, 5090-5098.

10 The DNA fragments may be obtained by digestion of DNA containing the required sequences of nucleotides with appropriate restriction enzymes, by chemical synthesis, by enzymatic polymerisation, or by a combination of these methods.

Digestion with restriction enzymes may be performed in an appropriate buffer at a temperature of 20°-70°C, generally in a volume of 50µl or less with 0.1-10µg DNA.

15 Enzymatic polymerisation of DNA may be carried out *in vitro* using a DNA polymerase such as DNA polymerase 1 (Klenow fragment) in an appropriate buffer containing the nucleoside triphosphates dATP, dCTP, dGTP and dTTP as required at a temperature of 10°-37°C, generally in a volume of 50µl or less.

20 Enzymatic ligation of DNA fragments may be carried out using a DNA ligase such as T4 DNA ligase in an appropriate buffer at a temperature of 4°C to 37°C, generally in a volume of 50µl or less.

25 The chemical synthesis of the DNA polymer or fragments may be carried out by conventional phosphotriester, phosphite or phosphoramidite chemistry, using solid phase techniques such as those described in 'Chemical and Enzymatic Synthesis of Gene
Fragments - A Laboratory Manual' (ed. H.G. Gassen and A. Lang), Verlag Chemie,
Weinheim (1982), or in other scientific publications, for example M.J.Gait, H.W.D.
Matthes M. Singh, B.S. Sproat and R.C. Titmas, *Nucleic Acids Research*, 1982, 10,
6243; B.S. Sproat and W. Bannwarth, *Tetrahedron Letters*, 1983, 24, 5771; M.D.
Matteucci and M.H. Caruthers, *Tetrahedron Letters*, 1980, 21, 719; M.D. Matteucci and
30 M.H. Caruthers, *Journal of the American Chemical Society*, 1981, 103, 3185; S.P.
Adams *et al.*, *Journal of the American Chemical Society*, 1983, 105, 661; N.D. Sinha, J.
Biernat, J. McMannus and H. Koester, *Nucleic Acids Research*, 1984, 12, 4539; and
H.W.D. Matthes *et al.*, *EMBO Journal*, 1984, 3, 801. Preferably an automated DNA
synthesiser (for example, Applied Biosystems 381A Synthesiser) is employed.

35 The DNA polymer is preferably prepared by ligating two or more DNA molecules which together comprise a DNA sequence encoding the polypeptide.

The DNA molecules may be obtained by the digestion with suitable restriction enzymes of vectors carrying the required coding sequences.

The precise structure of the DNA molecules and the way in which they are obtained depends upon the structure of the desired product. The design of a suitable
5 strategy for the construction of the DNA molecule coding for the polypeptide is a routine matter for the skilled worker in the art.

In particular, consideration may be given to the codon usage of the particular host cell. The codons may be optimised for high level expression in *E. coli* using the principles set out in Devereux *et al.*, (1984) Nucl. Acid Res., 12, 387.

10 The expression of the DNA polymer encoding the polypeptide in a recombinant host cell may be carried out by means of a replicable expression vector capable, in the host cell, of expressing the DNA polymer. Novel expression vectors also form part of the invention.

The replicable expression vector may be prepared in accordance with the
15 invention, by cleaving a vector compatible with the host cell to provide a linear DNA segment having an intact replicon, and combining said linear segment with one or more DNA molecules which, together with said linear segment, encode the polypeptide, under ligating conditions:

The ligation of the linear segment and more than one DNA molecule may be
20 carried out simultaneously or sequentially as desired.

Thus, the DNA polymer may be preformed or formed during the construction of the vector, as desired. The choice of vector will be determined in part by the host cell, which may be prokaryotic, such as *E. coli*, or eukaryotic, such as mouse C127, mouse myeloma, chinese hamster ovary, fungi e.g. filamentous fungi or unicellular 'yeast' or an
25 insect cell such as *Drosophila*. The host cell may also be in a transgenic animal. Suitable vectors include plasmids, bacteriophages, cosmids and recombinant viruses derived from, for example, baculoviruses or vaccinia.

The DNA polymer may be assembled into vectors designed for isolation of stable transformed mammalian cell lines expressing the fragment e.g. bovine papillomavirus
30 vectors in mouse C127 cells, or amplified vectors in chinese hamster ovary cells (DNA Cloning Vol. II D.M. Glover ed. IRL Press 1985; Kaufman, R.J. *et al.*. Molecular and Cellular Biology 5, 1750-1759, 1985; Pavlakis G.N. and Hamer, D.H. Proceedings of the National Academy of Sciences (USA) 80, 397-401, 1983; Goeddel, D.V. *et al.*, European Patent Application No. 0093619, 1983).

35 The preparation of the replicable expression vector may be carried out conventionally with appropriate enzymes for restriction, polymerisation and ligation of the DNA, by procedures described in, for example, Sambrook *et al.*, cited above.

Polymerisation and ligation may be performed as described above for the preparation of the DNA polymer. Digestion with restriction enzymes may be performed in an appropriate buffer at a temperature of 20°-70°C, generally in a volume of 50µl or less with 0.1-10µg DNA.

5 The recombinant host cell is prepared, in accordance with the invention, by transforming a host cell with a replicable expression vector of the invention under transforming conditions. Suitable transforming conditions are conventional and are described in, for example, Sambrook *et al.*, cited above, or "DNA Cloning" Vol. II, D.M. Glover ed., IRL Press Ltd, 1985.

10 The choice of transforming conditions is determined by the host cell. Thus, a bacterial host such as *E.coli*, may be treated with a solution of CaCl₂ (Cohen *et al.*, Proc. Nat. Acad. Sci., 1973, 69, 2110) or with a solution comprising a mixture of RbCl, MnCl₂, potassium acetate and glycerol, and then with 3-[N-morpholino]-propane-sulphonic acid, RbCl and glycerol or by electroporation as for example described by Bio-
15 Rad Laboratories, Richmond, California, USA, manufacturers of an electroporator. Mammalian cells in culture may be transformed by calcium co-precipitation of the vector DNA onto the cells or by using cationic liposomes.

 The invention also extends to a host cell transformed with a replicable expression vector of the invention.

20 Culturing the transformed host cell under conditions permitting expression of the DNA polymer is carried out conventionally, as described in, for example, Sambrook *et al.*, and "DNA Cloning" cited above. Thus, preferably the cell is supplied with nutrient and cultured at a temperature below 45°C.

 The protein product is recovered by conventional methods according to the host
25 cell. Thus, where the host cell is bacterial such as *E. coli* and the protein is expressed intracellularly, it may be lysed physically, chemically or enzymatically and the protein product isolated from the resulting lysate. Where the host cell is mammalian, the product is usually isolated from the nutrient medium.

 Where the host cell is bacterial, such as *E. coli*, the product obtained from the
30 culture may require folding for optimum functional activity. This is most likely if the protein is expressed as inclusion bodies. There are a number of aspects of the isolation and folding process that are regarded as important. In particular, the polypeptide is preferably partially purified before folding, in order to minimise formation of aggregates with contaminating proteins and minimise misfolding of the polypeptide. Thus, the
35 removal of contaminating *E. coli* proteins by specifically isolating the inclusion bodies and the subsequent additional purification prior to folding are important aspects of the procedure.

The folding process is carried out in such a way as to minimise aggregation of intermediate-folded states of the polypeptide. Thus, careful consideration needs to be given to, among others, the salt type and concentration, temperature, protein concentration, redox buffer concentrations and duration of folding. The exact condition
5 for any given polypeptide generally cannot be predicted and must be determined by experiment.

There are numerous methods available for the folding of proteins from inclusion bodies and these are known to the skilled worker in this field. The methods generally involve breaking all the disulphide bonds in the inclusion body, for example with 50mM
10 2-mercaptoethanol, in the presence of a high concentration of denaturant such as 8M urea or 6M guanidine hydrochloride. The next step is to remove these agents to allow folding of the proteins to occur. Formation of the disulphide bridges requires an oxidising environment and this may be provided in a number of ways, for example by air, or by incorporating a suitable redox system, for example a mixture of reduced and oxidised
15 glutathione.

Preferably, the inclusion body is solubilised using 8M urea, in the presence of mercaptoethanol, and protein is folded, after initial removal of contaminating proteins, by addition of cold buffer. Suitable buffers may be identified using the techniques described in I.Dodd *et al*, 'Perspectives in Protein Engineering and Complementary
20 Technologies', Mayflower Publications, 66-69, 1995. A suitable buffer for many of the SCR constructs described herein is 20mM ethanolamine containing 1mM reduced glutathione and 0.5mM oxidised glutathione. The folding is preferably carried out at a temperature in the range 1 to 50°C over a period of 1 to 4 days.

If any precipitation or aggregation is observed, the aggregated protein can be
25 removed in a number of ways, for example by centrifugation or by treatment with precipitants such as ammonium sulphate. Where either of these procedures are adopted, monomeric polypeptide is the major soluble product.

If the bacterial cell secretes the protein, folding is not usually necessary.

The polypeptide portion of the derivative of the invention may include a C-
30 terminal cysteine to facilitate post translational modification. A soluble polypeptide including a C-terminal cysteine also forms part of the invention. Expression in a bacterial system is preferred for proteins of moderate size (up to ~70kDa) and with <~8 disulphide bridges. More complex proteins for which a free terminal Cys could cause refolding or stability problems may require stable expression in mammalian cell lines
35 (especially CHO). This will also be needed if a carbohydrate membrane binding element is to be introduced post-translationally. The use of insect cells infected with recombinant baculovirus encoding the polypeptide portion is also a useful general method for

preparing more complex proteins and will be preferred when it is desired to carry out certain post-translational processes (such as palmitoylation) biosynthetically (see for example, M.J. Page *et al* J. Biol. Chem. 264, 19147-19154, 1989)

5 A preferred method of handling proteins C-terminally derivatised with cysteine is as a mixed disulphide with mercaptoethanol or glutathione or as the 2-nitro, 5-carboxyphenyl thio- derivative as generally described below in Methods.

Peptide membrane binding elements may be prepared using standard solid state synthesis such as the Merrifield method and this method can be adapted to incorporate required non-peptide membrane binding elements such as N-acyl groups derived from
10 myristic or palmitic acids at the N terminus of the peptide. In addition activation of an amino acid residue for subsequent linkage to a protein can be achieved during chemical synthesis of such membrane binding elements. Examples of such activations include formation of the mixed 2-pyridyl disulphide with a cysteine thiol or incorporation of an N-haloacetyl group. Both of these groups are capable of reaction with free thiols,
15 through disulphide interchange and alkylation, respectively. Peptides can optionally be prepared as the C-terminal amide and/or with a conventional N-terminal blocking group such as acetyl.

The invention also provides a peptidic membrane binding element comprising one or more derivatisations selected from:

- 20 a terminal cysteine residue optionally activated at the thiol group;
- an N-haloacetyl group (where halo signifies chlorine, bromine or iodine) located at the N-terminus of the peptide or at an ϵ -amino group of a lysine residue;
- an amide group at the C-terminus;
- an N-terminal blocking group; and
- 25 a fatty acid N-acyl group at the N-terminus or at an ϵ -amino group of a lysine residue.

Chemical bridging groups and reagents suitable for their formation include those described in EP0109653, EP0152736, EP0155388 and EP0284413, incorporated herein by reference. The bridging group is generally of the formula:

30 $-A-R-B-$ (I)

in which each of A and B, which may be the same or different, represents $-CO-$, $-C(=NH_2^+)-$, maleimido, $-S-$ or a bond and R is a bond or a linking group containing one or more $-(CH_2)-$ or meta-, ortho- or para- disubstituted phenyl units, preferably ortho or para, optionally together with a hydrophilic portion.

35 Where the polypeptide portion of the derivative of the invention and a peptidic membrane binding element both include a C-terminal cysteine the chemical bridging group will take the form $-S-S-$. The bridge is generated by conventional disulphide

exchange chemistry, by activating a thiol on one polypeptide and reacting the activated thiol with a free thiol on the other polypeptide. Such activation procedures make use of disulphides which form stable thiolate anions upon cleavage of the S-S linkage and include reagents such as 2,2' dithiopyridine and 5,5'-dithio(2-nitrobenzoic acid, DTNB) which form intermediate mixed disulphides capable of further reaction with thiols to give stable disulphide linkages.

R may include moieties which interact with water to maintain the water solubility of the linkage and suitable moieties include -CO-NH-, -CO-NMe-, -S-S-, -CH(OH)-, -SO₂-, -CO₂-, -(CH₂CH₂-O)_m- and -CH(COOH)- where m is an integer of 2 or more, or linear hydrophilic polymers such as polyethylene glycol, polypropylene glycol, polyglycine, polyalanine or polysarcosine.

Other examples of R include -(CH₂)_r-, -(CH₂)_p-S-S-(CH₂)_q- and -(CH₂)_p-CH(OH)-CH(OH)-(CH₂)_q-, in which r is an integer of at least 2, preferably at least 4 and p and q are independently integers of at least 2

In a further aspect R may take the form -U-V-W- where U is (CH₂)₂CONH(CH₂)_n in which n is an integer of 3 to 8, V is O, S, NR_a or NR_a-NR_a where each R_a is H or C₁₋₆ alkyl, NH-O or O-NH, and W is benzyl substituted at the 2- or 4- position by the group B. In a preferred embodiment R is (CH₂)₂CONH(CH₂)_nNH-(4-phenyl) where n is an integer of 3 to 8. The bridging group of formula (I) may be derived from a linking agent of formula (II):



in which R₁ is a bond or a linking group and X and Y are functional groups reactable with surface amino acid groups, preferably a lysine or cysteine group, the N-terminal amino group, a catalytic serine hydroxyl or a protein attachment group, and X, R₁- and Y are chosen so as to generate the required bridging group -A-R-B-.

Preferred agents are those where X and Y are different, known as heterobifunctional agents. Each end of the agent molecule is reacted in turn with each polypeptide to be linked in separate reactions. Examples of heterobifunctional agents of formula (II) include:

N-succinimidyl 3-(2-pyridyldithio) propionate
 succinimidyl 4-(N-maleimido) caproate
 3-(2-pyridyl) methyl propionimide hydrochloride
 4'-amidinophenyl 4-N-[2-N-(3-[2-pyridyldithio]ethylcarbonyl)aminoethyl]
 aminobenzoate hydrochloride.

Other suitable agents are disclosed in EP0109653, EP0152736, EP0155388 and EP0284413, in particular those of formula (II) in EP0155388 and (III) in EP0284413 incorporated herein by reference.

In each case Y is capable of reacting with a thiol group on a polypeptide, which may be a native thiol or one introduced as a protein attachment group.

The protein attachment group is a functionality derived by modification of a polypeptide or protein with a reagent specific for one or more amino acid side chains, and which contains a group capable of reacting with a cleavable section on the other polypeptide. An example of a protein attachment group is a thiol group. An example of a cleavable section is a disulphide bond. Alternatively the cleavable section may comprise an α , β dihydroxy function.

As an example, the introduction of a free thiol function by reaction of a polypeptide with 2-iminothiolane, N-succinimidyl 3-(2-pyridyldithio) propionate (with subsequent reduction) or N-acetyl homocysteine thiolactone will permit coupling of the protein attachment group with a thiol-reactive Y structure. Alternatively, the protein attachment group can contain a thiol-reactive entity such as the 6-maleimidoethyl group or a 2-pyridyl-dithio group which can react with a free thiol in X. Preferably, the protein attachment group is derived from protein modifying agents such as 2-iminothiolane that react with lysine ϵ -amino groups in proteins.

When X represents a group capable of reacting directly with the amino acid side chain of a protein, it is preferably an N-succinimidyl group. When X represents a group capable of reacting with a protein attachment group, it is preferably a pyridylthio group. When X represents a group capable of reacting with a catalytic serine hydroxyl it is preferably an 4-amidinophenyl ester group optionally substituted by one or more electron withdrawing groups which increases the reactivity of the ester, of the kind contained in the compounds of formula (II) in EP0155388 and (III) in EP0284413.

In the above processes, modification of a polypeptide to introduce a protein attachment group is preferably carried out in aqueous buffered media at a pH between 3.0 and 9.0 depending on the reagent used. For a preferred reagent, 2-iminothiolane, the pH is preferably 6.5-8.5. The concentration of polypeptide is preferably high (> 10mg/ml) and the modifying reagent is used in a moderate (1.1- to 5-fold) molar excess, depending on the reactivity of the reagent. The temperature and duration of reaction are preferably in the range 0°-40°C and 10 minutes to 7 days. The extent of modification of the polypeptide may be determined by assaying for attachment groups introduced.

Such assays may be standard protein chemical techniques such as titration with 5,5'-dithiobis-(2-nitrobenzoic acid). Preferably, 0.5-3.0 moles of protein attachment group will be introduced on average per mole of polypeptide. The modified polypeptide may be separated from excess modifying agents by standard techniques such as dialysis, ultrafiltration, gel filtration and solvent or salt precipitation. The intermediate material may be stored in frozen solution or lyophilised.

Where the linking agent of formula (II) contains an amidino phenyl ester group X the agent is preferably first reacted with a polypeptide enzyme via the group X and the reaction is preferably carried out under the conditions described for the introduction of blocking groups in European Published Patent Application No. 0,009,879. Having been
5 freed of excess reagent by suitable techniques such as high performance size exclusion chromatography or diafiltration, the acylated enzyme may then be reacted with the other polypeptide at a ratio of approximately 1:1 in a non-nucleophilic buffer at pH7.0-8.0 and 0°-30°C for up to 6h. However, it is preferable to conduct the coupling below 10°C (preferably 0°-4°C) in order to minimise the hydrolysis of the acylated enzyme.

10 Where a protein attachment group is introduced in this way, the bridging group (I) will be formed from a reaction of the linking agent (II) and the protein attachment group.

The polypeptides to be linked are reacted separately with the linking agent or the reagent for introducing a protein attachment group by typically adding an excess of the
15 reagent to the polypeptide, usually in a neutral or moderately alkaline buffer, and after reaction removing low molecular weight materials by gel filtration or dialysis. The precise conditions of pH, temperature, buffer and reaction time will depend on the nature of the reagent used and the polypeptide to be modified. The polypeptide linkage reaction is preferably carried out by mixing the modified polypeptides in neutral buffer in an
20 equimolar ratio. Other reaction conditions e.g. time and temperature, should be chosen to obtain the desired degree of linkage. If thiol exchange reactions are involved, the reaction should preferably be carried out under an atmosphere of nitrogen. Preferably, UV-active products are produced (eg from the release of pyridine 2-thione from 2-pyridyl dithio derivatives) so that coupling can be monitored.

25 After the linkage reaction, the polypeptide conjugate can be isolated by a number of chromatographic procedures such as gel filtration, ion-exchange chromatography, affinity chromatography or hydrophobic interaction chromatography. These procedures may be either low pressure or high performance variants.

The conjugate may be characterised by a number of techniques including low
30 pressure or high performance gel filtration, SDS polyacrylamide gel electrophoresis or isoelectric focussing.

Membrane binding elements which are fatty acid derivatives are attached post translationally to a peptidic membrane binding element, preferably at the terminus of the polypeptide chain. Preferably, where the recombinant polypeptide portion of the
35 derivative of the invention contains the peptidic membrane binding element, it has a unique cysteine for coupling to the fatty acid derivative. Where the recombinant polypeptide has a cysteine residue, a thiol-derivative of the fatty acid is added to the

refolded recombinant protein at a late stage in purification (but not necessarily the final stage) and at a reagent concentration preferably below the critical micelle concentration. One of the fatty acid derivative and the recombinant peptide will have the thiol group activated as described above for thiol interchange reactions. The fatty acid derivative is preferably a C₁₀₋₂₀ fatty acyl derivative of an aminoC₂₋₆alkane thiol (optionally C-substituted) such as N-(2-myristoyl) aminoethanethiol or N-myristoyl L-cysteine and forms part of the invention.

Suitable examples of hydrophilic synthetic polymers include polyethyleneglycol (PEG), preferably α,ω functionalised derivatives, more preferably α -amino , ω -carboxy-PEG of molecular weight between 400 and 5000 daltons which are linked to the polypeptide for example by solid-phase synthesis methods (amino group derivatisation) or by thiol-interchange chemistry.

Membrane binding elements derived from ligands of known integral membrane proteins, either amino acid sequences or carbohydrates, may be generated by post-translational modification using the glycosylation pathways of eukaryotic cells targeted to N-linked glycosylation sites in the peptide sequence.

Convenient generic final stage purification strategies are hydrophobic interaction chromatography (HIC) on C₂-C₈ media and cation exchange chromatography for separation of derivatised and underderivatised proteins into which a hydrophobic-electrostatic switch combination has been inserted.

The derivatives of this invention are preferably administered as pharmaceutical compositions.

Accordingly, the present invention also provides a pharmaceutical composition comprising a derivative of the invention in combination with a pharmaceutically acceptable carrier.

The compositions according to the invention may be formulated in accordance with routine procedures for administration by any route, such as oral, topical, parenteral, sublingual or transdermal or by inhalation. The compositions may be in the form of tablets, capsules, powders, granules, lozenges, creams or liquid preparations, such as oral or sterile parenteral solutions or suspensions or in the form of a spray, aerosol or other conventional method for inhalation.

The topical formulations of the present invention may be presented as, for instance, ointments, creams or lotions, eye ointments and eye or ear drops, impregnated dressings and aerosols, and may contain appropriate conventional additives such as preservatives, solvents to assist drug penetration and emollients in ointments and creams.

The formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may be

present as from about 1% up to about 98% of the formulation. More usually they will form up to about 80% of the formulation.

Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example
5 syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants, for example potato starch; or acceptable wetting agents such as sodium lauryl sulphate. Tablets may also contain agents for the stabilisation of polypeptide drugs against
10 proteolysis and absorption-enhancing agents for macromolecules. The tablets may be coated according to methods well known in normal pharmaceutical practice.

Suppositories will contain conventional suppository bases, e.g. cocoa-butter or other glyceride.

For parenteral administration, fluid unit dosage forms are prepared utilizing the
15 compound and a sterile vehicle, water being preferred. The compound, depending on the vehicle and concentration used, is dissolved in the vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilised before filling into a suitable vial or ampoule and sealing.

Parenteral formulations may include sustained-release systems such as
20 encapsulation within microspheres of biodegradable polymers such as poly-lactic co-glycolic acid.

Advantageously, agents such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. The dry
25 lyophilized powder is then sealed in the vial and an accompanying vial of water for injection may be supplied to reconstitute the liquid prior to use. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

Compositions of this invention may also suitably be presented for administration
30 to the respiratory tract as a snuff or an aerosol or solution for a nebulizer, or as a microfine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case the particles of active compound suitably have diameters of less than 50 microns, preferably less than 10 microns for example diameters in the range of 1-50 microns, 1-10 microns or 1-5 microns. Where appropriate, small amounts of
35 anti-asthmatics and bronchodilators, for example sympathomimetic amines such as isoprenaline, isoetharine, salbutamol, phenylephrine and ephedrine; xanthine derivatives

such as theophylline and aminophylline and corticosteroids such as prednisolone and adrenal stimulants such as ACTH may be included.

Microfine powder formulations may suitably be administered in an aerosol as a metered dose or by means of a suitable breath-activated device.

5 Suitable metered dose aerosol formulations comprise conventional propellants, cosolvents, such as ethanol, surfactants such as oleyl alcohol, lubricants such as oleyl alcohol, desiccants such as calcium sulphate and density modifiers such as sodium chloride.

10 Suitable solutions for a nebulizer are isotonic sterilised solutions, optionally buffered, at for example between pH 4-7, containing up to 20mg ml⁻¹ of compound but more generally 0.1 to 10mg ml⁻¹, for use with standard nebulisation equipment.

15 The quantity of material administered will depend upon the potency of the derivative and the nature of the complaint be decided according to the circumstances by the physician supervising treatment. However, in general, an effective amount of the polypeptide for the treatment of a disease or disorder is in the dose range of 0.01-100mg/kg per day, preferably 0.1mg-10mg/kg per day, administered in up to five doses or by infusion.

No adverse toxicological effects are indicated with the compounds of the invention within the above described dosage range.

20 The invention also provides a derivative of the invention for use as a medicament.

The invention further provides a method of treatment of disorders amenable to treatment by a soluble peptide which comprises administering a soluble derivative of said soluble peptide according to the invention, and the use of a derivative of the invention for the preparation of a medicament for treatment of such disorders.

25 In one preferred aspect the present invention relates to derivatives for use in the therapy of disorders involving complement activity and various inflammatory and immune disorders.

30 In this preferred aspect the soluble polypeptide which is derivatised in accordance with the invention is a soluble complement inhibitor such as a soluble CR1 polypeptide fragment.

35 Constituting about 10% of the globulins in normal serum, the complement system is composed of many different proteins that are important in the immune system's response to foreign antigens. The complement system becomes activated when its primary components are cleaved and the products alone or with other proteins, activate additional complement proteins resulting in a proteolytic cascade. Activation of the complement system leads to a variety of responses including increased vascular permeability, chemotaxis of phagocytic cells, activation of inflammatory cells,

opsonization of foreign particles, direct killing of cells and tissue damage. Activation of the complement system may be triggered by antigen-antibody complexes (the classical pathway) or, for example, by lipopolysaccharides present in cell walls of pathogenic bacteria (the alternative pathway).

5 Complement receptor type 1 (CR1) has been shown to be present on the membranes of erythrocytes, monocytes/macrophages, granulocytes, B cells, some T cells, splenic follicular dendritic cells, and glomerular podocytes. CR1 binds to the complement components C3b and C4b and has also been referred to as the C3b/C4b receptor. The structural organisation and primary sequence of one allotype of CR1 is
10 known (Klickstein *et al.*, 1987, J. Exp. Med. 165:1095-1112, Klickstein *et al.*, 1988, J. Exp. Med. 168:1699-1717; Hourcade *et al.*, 1988, J. Exp. Med. 168:1255-1270, WO 89/09220, WO 91/05047). It is composed of 30 short consensus repeats (SCRs) that each contain around 60-70 amino acids. In each SCR, around 29 of the average 65 amino acids are conserved. Each SCR has been proposed to form a three dimensional triple
15 loop structure through disulphide linkages with the third and first and the fourth and second half-cystines in disulphide bonds. CR1 is further arranged as 4 long homologous repeats (LHRs) of 7 SCRs each. Following a leader sequence, the CR1 molecule consists of the N-terminal LHR-A, the next two repeats, LHR-B and LHR-C, and the most C-terminal LHR-D followed by 2 additional SCRs, a 25 residue putative transmembrane
20 region and a 43 residue cytoplasmic tail.

Based on the mature CR1 molecule having a predicted N-terminal glutamine residue, hereinafter designated as residue 1, the first four SCR domains of LHR-A are defined herein as consisting of residues 2-58, 63-120, 125-191 and 197-252, respectively, of mature CR1.

25 Several soluble fragments of CR1 have been generated via recombinant DNA procedures by eliminating the transmembrane region from the DNAs being expressed (WO 89/09220, WO 91/05047). The soluble CR1 fragments were functionally active, bound C3b and/or C4b and demonstrated Factor I cofactor activity depending upon the regions they contained. Such constructs inhibited *in vitro* complement-related functions
30 such as neutrophil oxidative burst, complement mediated hemolysis, and C3a and C5a production. A particular soluble construct, sCR1/pBSCR1c, also demonstrated *in vivo* activity in a reversed passive Arthus reaction (WO 89/09220, WO 91/05047; Yeh *et al.*, 1991, J. Immunol. 146:250), suppressed post-ischemic myocardial inflammation and necrosis (WO 89/09220, WO 91/05047; Weisman *et al.*, Science, 1990, 249:146-1511;
35 Dupe, R. *et al.* Thrombosis & Haemostasis (1991) 65(6) 695.) and extended survival rates following transplantation (Pruitt & Bollinger, 1991, J. Surg. Res 50:350; Pruitt *et al.*, 1991 Transplantation 52; 868). Furthermore, co-formulation of sCR1/pBSCR1c


with p-anisoylated human plasminogen-streptokinase-activator complex (APSAC) resulted in similar anti-haemolytic activity as sCR1 alone, indicating that the combination of the complement inhibitor sCR1 with a thrombolytic agent was feasible (WO 91/05047).

- 5 The soluble CR1 polypeptide fragment encoded by sCR1/pBSCR1c, residues 1-1929 of CR1, may be derivatised in accordance with the invention.

Soluble polypeptides corresponding to part of CR1 have been found to possess functional complement inhibitory, including anti-haemolytic, activity. WO94/00571 discloses soluble polypeptides comprising, in sequence, one to four short consensus
10 repeats (SCR) selected from SCR 1, 2, 3 and 4 of long homologous repeat A (LHR-A) as the only structurally and functionally intact SCR domains of CR1 and including at least SCR3.

In preferred aspects, the polypeptide comprises, in sequence, SCR 1, 2, 3 and 4 of LHR-A or SCR 1, 2 and 3 of LHR-A as the only structurally and functionally intact
15 SCR domains of CR1.


In one aspect, the polypeptides may be represented symbolically as follows:



in which SCR1 represents residues 2-58 of mature CR1, SCR2 represents residues 63-120 of mature CR1, SCR3 represents residues 125-191 of mature CR1, and V¹, W¹,
20 X¹ and Y¹ represent bonds or short linking sequences of amino acids, preferably 1 to 5 residues in length and which are preferably derived from native interdomain sequences in CR1.

In a preferred embodiment of formula (I), W¹, X¹ and Y¹ represent residues 59-62, 121-124 and 192-196, respectively, of mature CR1 and V¹ represents residue 1 of mature CR1 optionally linked via its N-terminus to methionine.
25

In another aspect the polypeptides may be represented symbolically as follows:



in which SCR1, SCR2 and SCR3 are as hereinbefore defined, SCR4 represents residues 197-252 of mature CR1 and V², W², X², Y² and Z² represents bonds or short
30 linking sequences of amino acids, preferably 1 to 5 residues in length and which are preferably derived from native interdomain sequences in CR1.

In preferred embodiments of formula (II), W², X², Y² and Z² represent residues 59-62, 121-124, 192-196, and residues 253 respectively, of mature CR1 and V² represents residue 1 of mature CR1 optionally linked via its N-terminus to methionine.

35 In one particular embodiment of formula (B) arginine 235 is replaced by histidine.

In the preferred embodiment of formula (B), residue 235 is arginine.


In one further aspect, the polypeptide may be represented symbolically as follows:



in which SCR3 is as hereinbefore defined and X^3 and Y^3 represent bonds or short linking sequences of amino acids, preferably 1 to 5 residues in length and which are preferably derived from native interdomain sequences in CR1.

In a preferred embodiment of formula (C) X^3 represents amino acids 122-124 of mature CR1 optionally linked to methionine at its N-terminus and Y^4 represents amino acids 192-196 of mature CR1.

In another further aspect, the polypeptide may be represented symbolically as follows:



in which SCR3 and SCR4 are as hereinbefore defined and X^4 , Y^4 and Z^4 represent bonds or short linking sequences of amino acids, preferably 1 to 5 residues in length and which are preferably derived from native interdomain sequences in CR1.

In a preferred embodiment of formula (D) X^4 represents amino acids 122-124 of mature CR1 optionally linked to methionine at its N-terminus and Y^4 and Z^4 represent amino acids 192-196 and 253 respectively of mature CR1.

The soluble CR1 polypeptide is derivatised in accordance with the invention by any convenient strategy from those outlined above. In a preferred embodiment the soluble CR1 polypeptide consists of residues 1-196 of CR1 and with an N-terminal methionine and the derivative comprises a myristoyl group and one or more polypeptides sequence selected from

DGPKKKKKKSPSKSSGC
 GSSKSPSKKKKKKPGDC
 CDGPKKKKKKSPSKSSK
 SKDGKKKKKKSKTKC
 CSAAPSSGFRILLKLV
 AAPSVIGFRILLKVAGC
 and
 DGPSEILRGDFSSC
 (N-terminus on left).

The soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative of this invention is useful in the treatment of many complement-mediated or complement-related diseases and disorders including, but not limited to, those listed below.

Disease and Disorders Involving Complement**Neurological Disorders**

multiple sclerosis

stroke

5 Guillain Barré Syndrome

traumatic brain injury

Parkinson's disease

allergic encephalitis

Alzheimer's disease

10

Disorders of Inappropriate or Undesirable Complement Activation

haemodialysis complications

hyperacute allograft rejection

xenograft rejection

15 corneal graft rejection

interleukin-2 induced toxicity during IL-2 therapy

paroxysmal nocturnal haemoglobinuria

Inflammatory Disorders

20 inflammation of autoimmune diseases

Crohn's Disease

adult respiratory distress syndrome

thermal injury including burns or frostbite

uveitis

25 psoriasis

asthma

acute pancreatitis

Post-Ischemic Reperfusion Conditions

30 myocardial infarction

balloon angioplasty

atherosclerosis (cholesterol-induced) & restenosis

hypertension

post-pump syndrome in cardiopulmonary bypass or renal haemodialysis

35 renal ischemia

intestinal ischaemia

Infectious Diseases or Sepsis

multiple organ failure

septic shock

5 **Immune Complex Disorders and Autoimmune Diseases**

rheumatoid arthritis

systemic lupus erythematosus (SLE)

SLE nephritis

proliferative nephritis

10 glomerulonephritis

haemolytic anemia

myasthenia gravis

Reproductive Disorders

15 antibody- or complement-mediated infertility

Wound Healing

20 The present invention is also directed to a pharmaceutical composition for treating a disease or disorder associated with inflammation or inappropriate complement activation comprising a therapeutically effective amount of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative of this invention, and a pharmaceutically acceptable carrier or excipient.

25 The present invention also provides a method of treating a disease or disorder associated with inflammation or inappropriate complement activation comprising administering to a subject in need of such treatment a therapeutically effective amount of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative of this invention.

In the above methods, the subject is preferably a human.

30 Further provided is the use of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative of this invention in the manufacture of a medicament for the treatment of a disease or disorder associated with inflammation or inappropriate complement activation.

35 In order to inhibit complement activation and, at the same time, provide thrombolytic therapy, the present invention provides compositions which further comprise a therapeutically active amount of a thrombolytic agent. An effective amount of a thrombolytic agent is in the dose range of 0.01-10mg/kg; preferably 0.1-5mg/kg.

Preferred thrombolytic agents include, but are not limited to, streptokinase, human tissue type plasminogen activator and urokinase molecules and derivatives, fragments or conjugates thereof. The thrombolytic agents may comprise one or more chains that may be fused or reversibly linked to other agents to form hybrid molecules (EP-A-0297882 and EP 155387), such as, for example, urokinase linked to plasmin (EP-A-0152736), a fibrinolytic enzyme linked to a water-soluble polymer (EP-A-0183503). The thrombolytic agents may also comprise muteins of plasminogen activators (EP-A-0207589). In a preferred embodiment, the thrombolytic agent may comprise a reversibly blocked *in vitro* fibrinolytic enzyme as described in U.S. Patent No. 4,285,932. A most preferred enzyme is the p-anisoyl plasminogen-streptokinase activator complex, anistreplase as described in U.S. Patent No. 4,808,405 (Monk *et al.*, 1987, Drugs 34:25-49).

Routes of administration for the individual or combined therapeutic compositions of the present invention include standard routes, such as, for example, intravenous infusion or bolus injection. Active complement inhibitors and thrombolytic agents may be administered together or sequentially, in any order.

The present invention also provides a method for treating a thrombotic condition, in particular acute myocardial infarction, in a human or non-human animal. This method comprises administering to a human or animal in need of this treatment an effective amount of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative according to this invention and an effective amount of a thrombolytic agent.

Also provided is the use of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative of this invention and a thrombolytic agent in the manufacture of a medicament for the treatment of a thrombotic condition in a human or animal. Such methods and uses may be carried out as described in WO 91/05047.

This invention further provides a method for treating adult respiratory distress syndrome (ARDS) in a human or non-human animal. This method comprises administering to the patient an effective amount of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative according to this invention.

The invention also provides a method of delaying hyperacute allograft or hyperacute xenograft rejection in a human or non-human animal which receives a transplant by administering an effective amount of a soluble complement inhibitor, such as a soluble CR1 polypeptide, derivative according to this invention. Such administration may be to the patient or by application to the transplant prior to implantation.

The invention yet further provides a method of treating wounds in a human or non-human animal by administering by either topical or parenteral e.g. intravenous

routes, an effective amount of a soluble complement inhibitor, such as a soluble CR1 polypeptide derivative according to this invention.

5 In another preferred aspect the soluble polypeptide is a thrombolytic agent such as prourokinase, streptokinase, tissue-type plasminogen activator or reteplase and the derivative of the invention is useful in the treatment of thrombotic disorders such as acute myocardial infarction. The invention thus provides a pharmaceutical composition for treating thrombotic disorders comprising a therapeutically effective amount of a derivative of a thrombolytic agent according to the invention, and a pharmaceutically acceptable carrier or excipient. The invention further provides a method of treatment of
10 thrombotic disorders by administering an effective amount of a derivative of a thrombolytic agent according to the invention, and the use of such derivative in the preparation of a medicament for the treatment of thrombotic disorders.

The following Methods and Examples illustrate the invention.

GENERAL METHODS USED IN EXAMPLES

(i) DNA Cleavage

5 Cleavage of DNA by restriction endonucleases was carried out according to the manufacturer's instructions using supplied buffers. Double digests were carried out simultaneously if the buffer conditions were suitable for both enzymes. Otherwise double digests were carried out sequentially where the enzyme requiring the lowest salt condition was added first to the digest. Once the digest was complete the salt concentration was altered and the second enzyme added.

10

(ii) DNA ligation

Ligations were carried out using T4 DNA ligase purchased from Promega, as described in Sambrook *et al.*, (1989) Molecular Cloning: A Laboratory Manual 2nd Edition. Cold Spring Harbour Laboratory Press.

15

(iii) Plasmid isolation

Plasmid isolation was carried out by the alkaline lysis method described in Sambrook *et al.*, (1989) Molecular Cloning: A Laboratory Manual 2nd Edition. Cold Spring Harbour Laboratory Press or by one of two commercially available kits: the Promega WizardTM Plus Minipreps or Qiagen Plasmid Maxi kit according to the manufacturer's instructions.

20

(iv) DNA fragment isolation

DNA fragments were excised from agarose gels and DNA extracted using one of three commercially available kits: the QIAEX gel extraction kit or Qiaquick gel extraction kit (QIAGEN Inc., USA), or GeneClean (Bio 101 Inc, USA) according to the manufacturer's instructions.

25

(v) Introduction of DNA into *E. coli*

30 Plasmids were transformed into *E. coli* BL21(DE3) (Studier and Moffat, (1986), *J. Mol. Biol* 189:113), *E. coli* XLI-blue, BL21 (DE3) pLys-S or BLR (DE3) pLys-S that had been made competent using calcium chloride as described in Sambrook *et al.*, (1989). Cell lines were purchased as frozen competent cultures from Stratagene. *E. coli* JM109 was purchased as a frozen competent culture from Promega.

35

(vi) **DNA sequencing**

Plasmid DNA was sequenced on a Vistra DNA Labstation 625. The sequencing chemistry was performed using Amersham International's Thermo Sequenase fluorescent dye-terminator cycle sequencing kit' (RPN 2435), in conjunction with their TFP

- 5 fluorescent dye-terminator precipitation kit' (RPN 2433) according to the manufacturer's instructions.

The sequences produced by the above procedure were analysed by a Perkin Elmer ABI Prism 377 DNA Sequencer. This is an electrophoretic technique using 36 cm x 0.2mm 4% acrylamide gels, the fluorescently labeled DNA fragments being detected by a charge coupled device camera according to the manufacturer's instructions.

(vii) **Production of oligonucleotides**

Oligonucleotides were purchased from Cruachem.

15 (viii) **pBROC413**

The plasmid pT7-7 [Tabor, S (1990), Current Protocols in Molecular Biology, F. A. Ausubel, Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl,eds.] pp.16.2.1-16.2.11, Greene Publishing and Wiley-Interscience, New York.] contains DNA corresponding to nucleotides 2065-4362 of pBR322 and like pBR322 can be mobilized by a conjugative plasmid in the presence of a third plasmid ColK. A mobility protein encoded by ColK acts on the *nic* site at nucleotide 2254 of pBR322 initiating mobilization from this point. pT7-7 was digested with *LspI* and *BglII* and the protruding 5' ends filled in with the Klenow fragment of DNA Polymerase I. The plasmid DNA fragment was purified by agarose gel electrophoresis, the blunt ends ligated together and transformed into *E. coli* DH1 by electroporation using a Bio-Rad Gene Pulser and following the manufacturers recommended conditions. The resultant plasmid pBROC413 was identified by restriction enzyme analysis of plasmid DNA.

The deletion in pBROC413 from the *LspI* site immediately upstream of the *f* 10 promoter to the *BglII* site at nucleotide 434 of pT7-7 deletes the DNA corresponding to nucleotides 2065-2297 of pBR322. The *nic* site and adjacent sequences are therefore deleted making pBROC413 non mobilizable.

(ix) **Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS PAGE)**

SDS PAGE was carried out generally using the Novex system (British Biotechnology) according to the manufacturer's instructions. Prepacked gels of 4 - 20% acrylamide were used. Samples for electrophoresis, including protein molecular weight standards (for example LMW Kit, Pharmacia or Novex Mark 12) were usually diluted in

1%(w/v)SDS - containing buffer (with or without 5%(v/v) 2-mercaptoethanol), and left at room temperature for about 10 to 30 min before application to the gel.

(x) **Reduction of disulphides and modification of thiols in proteins**

5 There are a number of methods used for achieving the title goals. The reason it may be necessary to carry out selective reduction of disulphides is that during the isolation and purification of multi-thiol proteins, in particular during refolding of fully denatured multi-thiol proteins, inappropriate disulphide pairing can occur. In addition, even if correct disulphide pairing does occur, it is possible that a free cysteine in the
10 protein may become blocked, for example with glutathione. These derivatives are generally quite stable. In order to make them more reactive, for example for subsequent conjugation to another functional group, they need to be selectively reduced, with for example dithiothreitol (DTT) or Tris (2-carboxyethyl) phosphine.HCl (TCEP) then optionally modified with a function which is moderately unstable. An example of the
15 latter is Ellmans reagent (DTNB) which gives a mixed disulphide. In the case where treatment with DTNB is omitted, careful attention to experimental design is necessary to ensure that dimerisation of the free thiol-containing protein is minimised. Reference to the term 'selectively reduced' above means that reaction conditions eg. duration, temperature, molar ratios of reactants have to be carefully controlled so that reduction of
20 disulphide bridges within the natural architecture of the protein is minimised. All the reagents are commercially available eg. from Sigma or Pierce.

 The following general examples illustrate the type of conditions that may be used and that are useful for the generation of free thiols and their optional modification. The specific reaction conditions to achieve optimal thiol reduction and/or modification are
25 ideally determined for each protein batch.

 TCEP may be prepared as a 20mM solution in 50mM Hepes (approx. pH 4.5) and may be stored at -40 degrees C. DTT may be prepared at 10mM in sodium phosphate pH 7.0 and may be stored at -40 degrees C. DTNB may be prepared at 10mM in sodium phosphate pH 7.0 and may be stored at -40 degrees C. All of the above reagents are
30 typically used at molar equivalence or molar excess over protein concentration, the precise concentrations ideally identified experimentally. The duration and the temperature of the reaction are similarly determined experimentally. Generally the duration would be in the range 1 to 24 hours and the temperature would be in the range 2 to 30 degrees C. Excess reagent may be conveniently removed by buffer exchange, for
35 example using Sephadex G25 or Sephadex G50. A suitable buffer is 0.1M sodium phosphate pH7.0 or the solution may be left untreated.

EXAMPLES

Example 1 Preparation of N-(Myristoyl) 2-aminoethane thiol (MAET)

Myristoyl chloride (1.0 mmol) was added with vigorous stirring to ice-cooled dry pyridine (1.0 ml), and followed immediately by N-hydroxysuccinimide (1.5 mmol). The mixture was stirred for 4 h at ambient temperature (~23°C). 2-aminoethanethiol free base (1.1 mmol) was added as solid to the mixture and allowed to react for 6 h at ambient temperature, followed by 3 days at 4°C. The product was treated with water (5 ml), stirred for 1 h at ambient and filtered, washing with cold water. The white solid was dissolved in dimethylsulphoxide and reprecipitated with water and then vacuum dried over phosphorous pentoxide. The final yield was 0.21 g (~70%). Thiol titration using Ellman's reagent indicated that the product contained ~45% free thiol.

Example 2 Synthesis of Myristoyl/Electrostatic Switch Peptide Reagent 1 (MSWP-1) (SEQ ID NO: 27)

N-(Myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Pro-Gly-Asp-(S-2-Thiopyridyl)Cys-NH₂

The peptide:

Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Pro-Gly-Asp-Cys-NH₂
(SEQ ID NO: 5)

was prepared using solid phase synthesis via the general Fmoc/tBu strategy developed by Sheppard and Atherton (E. Atherton and R.C. Sheppard, Solid Phase Synthesis, IRL Press, Oxford, 1989). Kieselguhr-supported polydimethylacrylamide resin (Macrosorb 100) was used as the solid support and was derivatised with ethylene diamine.

Coupling reactions were carried out using N-α-Fmoc protected reagents pre-activated with N,N'-diisopropylcarbodiimide/ N-hydroxybenzotriazole (in 4-fold molar excess) with bromophenol blue monitoring. Fmoc cleavages used 20% piperidine in DMF. Reactions to assemble the peptide chain were carried out by repeated cycles of coupling and deprotection including the attachment of the modified Rink linkage reagent (p-[(R,S)-α-[1-(9H-fluorenyl-9-yl)-methoxyformamido] 2,4 dimethoxybenzyl]-phenoxyacetic acid) designed to yield a C-terminal amide on final cleavage. The side chain functionalities of the individual amino-acids were protected as follows:

Ser (tButyl), Lys (Boc), Asp (O-tButyl), Cys (Trityl).

On completion of the peptide assembly and with the peptide still attached to the resin, the myristoyl group was attached to the amino group of the N terminal glycine by direct coupling of myristic acid by the same activation procedure. This modified peptide was then cleaved from the resin and the side-chain protecting groups removed at the

same time by treatment with trifluoroacetic acid containing 2.5% water and 2.5% triisopropyl silane.

The crude product was treated with 2,2' dithiopyridine in 0.01M ammonium acetate solution at pH 8-9 for approx. 2h, then acidified with acetic acid and purified by preparative high performance liquid chromatography (HPLC) in 0.1% trifluoroacetic acid (TFA) /water and 0.1% TFA/acetonitrile as gradient component. After lyophilisation, the peptide was a white amorphous powder, soluble to at least 10mg/ml in dimethylsulphoxide. Fast atom bombardment mass spectrometry gave main peaks at m/e 2107.8, 2129.7 and 2145.8, corresponding to the monoprotated, monosodiated and monopotassiated molecular ions of the peptide. The 2-thiopyridyl content of the peptide was measured by dissolving it to around 0.03mM to 0.2 mM in 0.1M Sodium Borate pH 8.0 and reducing by addition of dithiothreitol to 5mM. The change in optical density at 343nm was used to calculate the amount of pyridine 2-thione released using an extinction coefficient at this wavelength of $8080 \text{ cm}^{-1} \text{ M}^{-1}$. This indicated that the peptide content was approximately 60% of the dry weight.

Example 3 Synthesis of Myristoyl/Electrostatic Switch Peptide Reagent 2 (MSWP-2) (SEQ ID NO: 28)

N-acetyl-Cys (2-thiopyridyl) Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Pro-Ser Lys-Ser-Ser-Lys-
(ϵ -N-(Myristoyl))-NH₂

The peptide:

Cys-Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Pro-Ser-Lys-Ser-Ser-Lys-NH₂ (SEQ ID NO: 18) was prepared by solid-phase synthesis using the general method described in Example 2 and with the following variations:

- a. The C-terminal lysine was protected by alkylation with the 4-methyl trityl (MTT) group; all other lysines were N- ϵ protected with the t-Boc group
- b. MTT was removed with 1% v/v trifluoroacetic acid in dichloromethane and the resulting unique free amino group derivatised with myristic acid prior to deprotection of the other lysines (as described in Example 2)

The N-terminus was acetylated with acetic anhydride upon completion of the peptide chain assembly. Generation of the 2-pyridyldithiocysteine moiety was by reaction of the deprotected peptide with 2,2'-dithiopyridine as described above. The product was purified as described in Example 2. Fast-atom bombardment mass spectrometry gave a molecular ion peak at 2221.3 (cf 2220.3 for the monoprotated theoretical mass).

Amino-acid Analysis:

	Asx	Ser	Gly	Pro
Theory:	1.0	4.0	1.0	2.0
Found	0.97	3.53	1.15	1.88

5 (Asx = Asn or Asp)

Amino-acid analysis indicated a net peptide content by weight of 68.7%. The 2-pyridyl disulphide content was approximately 60% by weight using the method of Example 2.

Example 4 Synthesis of Myristoyl/Electrostatic Switch Peptide Reagent 3

10 (MSWP-3) (SEQ ID NO: 29)

N-(Myristoyl)-Ser-Lys-Asp-Gly-Lys-Lys-Lys-Lys-Lys-Ser-Lys-Thr-Lys-(S-2-Thiopyridyl)Cys-NH₂

The peptide:

15 Ser-Lys-Asp-Gly-Lys-Lys-Lys-Lys-Lys-Ser-Lys-Thr-Lys-Cys (SEQ ID NO: 19) was prepared using the general solid-phase synthesis protocol of Example 2. Myristoylation, C-terminal amidation and derivatisation of the Cys residue were performed as described in Example 2. After purification, mass spectrometry gave the major peak at 2040.5, corresponding to a monoprotonated form (Theory: 2039.5)

20

Amino-acid analysis:

	Asx	Ser	Gly	Thr	Lys
Theory:	1	2	1	1	9
Found:	1.02	2.04	1.14	1.06	8.85

25

The peptide content was about 56% by weight

Example 5 Synthesis of T-Cell Targeting Peptide Reagent 1 (TCTP-1) (SEQ ID NO: 30)

30 N-acetyl-(2-thiopyridyl)Cys Ser-Ala-Ala-Pro-Ser-Ser-Gly-Phe-Arg-Ile-Leu-Leu-Lys-Val-CONH(CH₂)₉CH₃

The peptide Cys-Ser-Ala-Ala-Pro-Ser-Ser-Gly-Phe-Arg-Ile-Leu-Leu-Lys-Val (SEQ ID NO: 20) was prepared using the general solid-phase methodology of Example 2 and N-acetylated as in Example 3. The C-terminus was derivatised using n-decylamine in place of the Rink reagent. Mass spectrometry of the purified peptide gave a major peak at 1952.3 corresponding to a monoprotonated molecular ion (Theory:

35

1951.1.) An ion at 1843.3 was also observed, this is believed to correspond to loss of the thiopyridyl group in the spectrophotometer.

Amino-acid analysis:

5		Ser	Gly	Arg	Ala	Pro	Val	Ile	Phe	Leu	Lys
	Theory:	3	1	1	2	1	1	1	1	3	1
	Found:	2.95	1.10	1.10	2.11	1.04	0.60	0.92	1.00	3.03	1.03

The peptide content by weight was 53%

10

Example 6 Expression and isolation of [SCR1-3]-Cys (SEQ ID NO: 6)

(a) Construction of plasmid pDB1030 encoding [SCR 1-3]-Cys

The plasmid coding for SCR1-3 of LHR-A of CR1, pDB1013-5 (patent application WO 94/00571) was digested with restriction endonucleases *EcoRI* and *HindIII* and the 2.2 kB plasmid band was isolated from an agarose gel using a Qiagen Qiaex DNA extraction kit according to the manufacturer's instructions. This is fragment 1. A second batch of pDB1013-5 was digested with *BanI* and *EcoRI* and the 196 bp band was extracted from agarose as above. This is fragment 2. Two oligonucleotides, SEQ ID No.1 and SEQ ID No.2, were annealed to give a final DNA concentration of 100 pmoles/ul. The annealed oligo has a *BanI/EcoRI* overhang and duplicates the sequence at the 3' end of pDB1013-5 but in addition contains a codon coding for cysteine just before the stop codon. This is fragment 3.

Fragments 1,2 and 3 were ligated with T4 DNA ligase in a single reaction to give pDB1030. The ligated plasmid was transformed into competent *E. coli* JM109 purchased from Promega. Resulting colonies were analysed by restriction endonuclease digestion and DNA sequencing confirmed that the encoded amino acid sequence of SCR(1-3) (SEQ ID No.27 of WO 94/00571) had been altered by a single C-terminal cysteine residue to give SEQ ID No.6.

30 (b) Expression of [SCR1-3]-Cys from pDB1030

pDB1030 was transformed into calcium chloride competent *E. coli* BL21(DE3) and resultant colonies were isolated and checked for plasmid content. To express protein from pDB1030 in *E. coli* BL21(DE3), a single colony was inoculated into 10 ml LB-phosphate media (20g/L tryptone, 15g/L yeast extract, 0.8g/L NaCl, 0.2g/L Na₂HPO₄, 0.1g/L KH₂PO₄) containing 50ug/ml ampicillin. The culture was grown for 6 hours at 37°C, 230 r.p.m. before being used to inoculate 100 ml of the same media containing 50 ug/ml ampicillin. Growth was under the same conditions overnight. 25 ml of each

culture were then used to inoculate 600 ml of the same media with 50 ug/ml ampicillin in 3 L erlenmeyer flasks. Cells were grown to an OD of 0.8 - 1.0 at A₆₀₀ nm. IPTG (isopropyl B-D galactopyranoside) was added to a final concentration of 1 mM and cells allowed to continue growth for a further 3 - 4 hours before harvesting by centrifugation
5 at 8000 g/10 min. Pellet from 2L of culture was stored at -80°C

(c) **Isolation, refolding, purification and formulation of [SCR1-3]-Cys**

The methods described are essentially those detailed in Dodd I. et al (1995) Protein Expression and Purification 6 727-736.

10 i) **Isolation of solubilised inclusion bodies**

The frozen cell pellet of *E. coli* BL21(DE3) (pDB1030) was resuspended in 50 mM Tris/50 mM NaCl/1 mM EDTA/0.1 mM PMSF pH 8.0 at a ratio of 33 ml for each litre of culture pellet. The suspension was transferred to a glass beaker surrounded by ice and sonicated (Heat systems - Ultrasonics W380; 50 x 50% pulse, pulse time = 5 sec.)
15 for typically 3 - 6 minutes. The disrupted pellet was then frozen and stored at -80°C. Approx. 2 weeks later the sonicate was thawed and centrifuged at approx. 8000g for 20 min. The pellet was resuspended in 20 mM Tris/8M urea/1 mM EDTA/50 mM 2-mercaptoethanol pH 8.5 (200ml) at room temperature by vigorous swirling, then left for 1h at room temperature followed by overnight at 4° C.

20 ii) **Initial purification using SP-Sepharose**

To the viscous solution was added SP-Sepharose FF (approx. 30g wet weight) that had been water washed and suction-dried. The mixture was swirled vigorously and left static for 1-2h at room temperature. The supernatant was decanted, sampled and discarded. The remaining slurry was resuspended to a uniform suspension and poured
25 into a glass jacket and allowed to settle into a packed bed. The column was equilibrated with 0.02M Tris/8M urea/0.05M 2-mercaptoethanol/0.001 M EDTA pH 8.5.at 4°C. When the A₂₈₀ of the eluate had stabilised at baseline, the buffer was changed to equilibration buffer additionally containing 1M NaCl. A single A₂₈₀ peak was eluted by the 1M NaCl-containing buffer; the volume was approx. 50ml. The protein
30 concentration of the solution was estimated by A₂₈₀ determination, using a molar extinction coefficient of 25000 cm⁻¹ of a sample that had been buffer-exchanged (Sephadex G25) into 50 mM formic acid. This showed the product had a protein concentration of 1.6mg/ml. The solution was stored at -40°C.

iii) **Folding and further processing**

35 25ml of the SP-Sepharose-purified product was added gradually over a 1 min period to 780 ml freshly prepared, cold 0.02M ethanolamine/1mM EDTA with continuous swirling, and left static for 1 h/40°C. Reduced glutathione (GSH) was added

to 1 mM and oxidised glutathione (GSSG) was added to 0.5 mM. The solution was clear and was left static approx 2-3°C for 3 d. The solution was then ultrafiltered using a YM10 membrane to a final retentate volume of about 35 ml; the retentate was slightly cloudy and had the appearance of a translucent solution. It was stored for 12 days at 4° C. It was then spun at 30 000g for 15 mins and the supernatant mixed with 9 vol. 0.1M NaH₂PO₄/1M (NH₄)₂SO₄ pH 7.0 (Buffer A) at room temperature and immediately centrifuged at 3000 rpm for 15 min. The supernatant was ultrafiltered (YM10) to about 4ml and then buffer-exchanged into 0.1M sodium phosphate pH 7.0 (5.0ml); this solution had a protein concentration of 1.7mg/ml by A280 analysis. It was treated with dithio bis nitrobenzoic acid (DTNB) (8-fold molar excess) for 30 min at room temperature. Free thiol content based on A412 measurement and an extinction coefficient (for the free thionitrobenzoate ion) of 13 600 was 6uM equivalent to only about 10% derivatisation to give Product A. The majority of the product was believed to be [SCR1-3]-Cys where the free C-terminal thiol was blocked by reaction with glutathione or 2-mercaptoethanol during the refolding stage.

(d) Alternative method for isolation, refolding, purification and formulation of [SCR1-3]-Cys

The method was similar to that described above, except that it more closely followed the procedures described in Dodd et al (op cit.). Notably, the ultrafiltered retentate post refolding was immediately treated with ammonium sulphate followed by clarification by centrifugation and Butyl Toyopearl chromatography. The resulting A280-absorbing fractions that eluted at about 0.2 to 0.4M ammonium sulphate were pooled and regarded as Product B. Starting with a nominal 100mg of fully reduced SCR1-3/cys, Product B contained 17mg. The product contained one major species by non-reduced SDS PAGE with an estimated purity of >90% and an apparent molecular weight of 21 000. On the basis of studies with similarly produced preparations it was believed to be the S-glutathione and/or S-mercaptoethanol derivatised form of the parent protein, although at least some batches produced in a similar way or stored for a period of time might exist as the free cysteine variant. The product also contained a polypeptide with an apparent molecular weight of about 40 000. On the basis of studies with similar batches of protein enriched in this species it was identified as the dimer of [SCR1-3]-Cys.

Example 7 Expression and isolation of SCR1-3/switch fusion (SEQ ID NO: 7)

H₂N-[SCR 1-3]-Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Ser-Pro-Ser-Lys-Ser-Ser-Gly-Cys-OH

(a) **Construction of plasmid pDB1031 encoding SCR1-3/switch**

Fragment 1 and fragment 2 of pDB1013-5 were the same as Example 6 above.

Two oligonucleotides, SEQ ID No.3 and SEQ ID No. 4, prepared by Cruachem were annealed to give a final DNA concentration of 100 pmoles/ul. The annealed oligo has an
5 *BanI/EcoRI* overhang and duplicates the sequence at the 3' end of pDB1013-5 but in addition contains 17 additional codons coding for DGPKKKKKKSPSKSSGC just before the stop codon. This is fragment 4.

Fragments 1, 2 and 4 were ligated with T4 DNA ligase in a single reaction to give pDB1031. The ligated plasmid was transformed into competent *E. coli* JM109.

- 10 Resulting colonies were analysed by restriction endonuclease digestion and DNA sequencing confirmed that the encoded amino acid sequence of SCR1-3 (SEQ ID 27 of WO 94/00571) had been altered by C terminal addition of amino acids DGPKKKKKKSPSKSSGC to give SEQ ID NO: 7.

15 (b) **Expression of SCR1-3/switch from pDB1031**

pDB1031 was transformed into calcium chloride competent *E. coli* BL21(DE3) and resultant colonies were isolated and checked for plasmid content. To express protein from pDB1031 in *E. coli* BL21(DE3), a single colony was inoculated into 10 ml LB-phosphate media (20g/L tryptone, 15g/L yeast extract, 0.8g/L NaCl, 0.2g/L

- 20 Na₂HPO₄, 0.1g/L KH₂PO₄) containing 50ug/ml ampicillin. The culture was grown for 6 hours at 37°C, 230 r.p.m. before being used to inoculate 100 ml of the same media containing 50 ug/ml ampicillin. Growth was under the same conditions overnight. 25 ml of each culture were then used to inoculate 600 ml of the same media with 50 ug/ml ampicillin in 3 L erlenmeyer flasks. Cells were grown to an OD of 0.8 - 1.0 at A₆₀₀ nm.
25 IPTG (isopropyl B-D galactopyranoside) was added to a final concentration of 1 mM and cells allowed to continue growth for a further 3 - 4 hours before harvesting by centrifugation at 8000 g/10 min. The cell pellet was frozen at -40 degrees C.

(c) **Isolation, refolding, purification and formulation of SCR1-3/switch**

- 30 The methods described are essentially those detailed in Dodd I. et al (1995) Protein Expression and Purification 6 727-736, with some modifications.

i) **Isolation of solubilised inclusion bodies**

- The frozen cell pellet of *E. coli* BL21(DE3) (pDB1031) was thawed and resuspended in 50 mM Tris/50 mM NaCl/1 mM EDTA/0.1 mM PMSF pH 8.0 at a ratio
35 of 33 ml for each litre of culture pellet. The suspension was transferred to a glass beaker surrounded by ice and sonicated (Heat systems - Ultrasonics W380; 50 x 50% pulse, pulse time = 5 sec.) for typically 3 - 6 minutes. The disrupted pellet was then frozen and

stored at -80°C. Approx. 1d later the sonicate was thawed and centrifuged at approx. 8000g for 20 min. The pellet was resuspended in 20 mM Tris/8M urea/1 mM EDTA/50 mM 2-mercaptoethanol pH 8.5 (240ml) at room temperature by vigorous swirling, then left for 1h at room temperature followed by 5 days at 4° C.

5 ii) **Preliminary purification using SP-Sepharose**

To the viscous solution was added SP-Sepharose FF (approx. 30g wet weight) that had been water washed and suction dried. The mixture was swirled vigorously and left static for approx. 2h at room temperature. The supernatant was decanted, sampled and discarded. The remaining slurry was resuspended to a uniform suspension and
10 poured into a glass jacket and allowed to settle into a packed bed. The column was equilibrated with 0.02M Tris/8M urea/0.05M 2-mercaptoethanol/0.001 M EDTA pH 8.5.at 4°C. When the A₂₈₀ of the eluate had stabilised at baseline, the buffer was changed to equilibration buffer additionally containing 1M NaCl. A single A₂₈₀ peak was eluted by the 1M NaCl-containing buffer; the volume was approx. 50ml. The
15 protein concentration of the solution was estimated by A₂₈₀ determination, using a molar extinction coefficient of 25000 cm⁻¹, of a sample that had been buffer-exchanged (Sephadex G25) into 50 mM formic acid. This showed the product had a protein concentration of 2.8mg/ml. Analysis by SDS PAGE/stain showed a major band (approx 80%) at about 23 000Da. The solution was stored at -40°C.

20 iii) **Folding and further processing**

14ml of the SP-Sepharose-purified product was added gradually over a 1 min period to 430 ml freshly prepared, cold 0.05M Hepes/2 M sodium chloride/1mM EDTA pH 8.0 with continuous swirling, and left static for 1 h/4°C. Reduced glutathione (GSH) was added to 1 mM and oxidised glutathione (GSSG) was added to 0.5 mM. The
25 solution was clear and was left static approx 2-3°C for 3 d. The solution was then ultrafiltered using a YM10 membrane to a final retentate volume of about 34 ml; the retentate was slightly cloudy. It was then spun at 25 000g for 15 mins and the supernatant buffer-exchanged into 0.1M sodium phosphate pH 7.0 (46 ml). This fraction contained 2 mg of protein on the basis of an A₂₈₀ determination. The solution
30 was mixed with DTNB (20mM; 0.65ml) for 20 min at 4 degrees C and then ultrafiltered to 2.4ml. This retentate was buffer-exchanged into 0.1M sodium phosphate pH 7.0 (3.0ml) and stored at -40 degrees C. Absorbance measurements at 412nm on the solution prior to ultrafiltration suggested 25% derivatisation with DTNB.

(d) Alternative isolation, refolding, purification and formulation of SCR1-3/switch

The method was similar to that described in (c) above, except that following the ultrafiltration step after refolding it more closely followed the procedures described in Dodd et al (*op cit.*). Notably, the ultrafiltered retentate post refolding was immediately treated with ammonium sulphate followed by clarification by centrifugation and Butyl Sepharose chromatography. The resulting A280-absorbing fractions that eluted at about 0.2 to 0.4M ammonium sulphate were pooled and regarded as initial product. Additional treatment with TCEP essentially as above, followed by DTNB yielded a final product at 10uM final protein concentration. The final product contained one major species by non-reduced SDS PAGE with an estimated purity of >90% and an apparent molecular weight of 23 000 and contained about 2 moles TNB per mole of protein.

Example 8 Preparation of [SCR1-3]-Cys-S-S-[MSWP-1] (SEQ ID NO: 8)




(a) Product A of Example 6(c) (1.5ml) was treated with dithiothreitol (30 ul of 0.5M in water, final concentration 10mM) for 60 min at 4°C to give the free peptide SEQ ID NO 6. The yellow solution was gel filtered at 4°C on a small column of Sephadex G-25 (PD-10, Pharmacia) into 0.05M Hepes.HCl buffer pH 7.5 (3.0ml). The slightly cloudy solution was mixed with a solution of MSWP-1 (Example 2) (3.8mM dithiopyridyl equivalents, 150 ul) to a final concentration of 0.18mM (~8 molar equivalents). The mixture was held for 2h on ice and then gel filtered as before but using 2 PD10 columns (1.6 ml applied, 3.2ml eluted). The final eluate was not cloudy and was stored frozen at -70°C in aliquots of 0.4ml.

(b) [SCR1-3]-Cys protein product B described in Example 6(d) (1.5ml; 31uM protein) was mixed with TCEP (20mM; 0.007ml) and incubated at room temperature for 23 h to give the free protein SEQ ID NO: 6. MSWP-1 (Example 2) (10mM; 0.093ml) was added and the solution incubated for a further 4 h. 0.75ml of the final solution was buffer-exchanged into 50mM formic acid and aliquots left in solution or lyophilised. The product was >80% pure by SDS PAGE and had an apparent molecular weight of 23 000, clearly shifted from the original parent molecular weight of 21 000. The

lyophilisate was easily soluble in 50mM formic acid at an estimated protein concentration of 2 mg/ml.

- (c) [SCR1-3]-Cys protein product B described in Example 6(d) (21.6ml; 31uM protein) was mixed with TCEP (20mM; 0.1ml) and incubated at room temperature for 22h to give the free protein SEQ ID NO 6. MSWP-1 (20mM in 0.1M sodium phosphate pH 7.0; 0.67ml) was added and the solution incubated for a further 4 h. All 22ml was buffer-exchanged into 50mM formic acid using Sephadex G50 (Vt 160ml). Three A280 peaks were obtained. The first one, eluting at volume 56 -106ml, was the title compound according to SDS PAGE analysis. The fraction was aliquoted and aliquots stored at - 40 degrees C or lyophilised. Amino acid analysis of the pre-lyophilisation solution indicated a protein concentration of 0.42mg/ml. A280 (1cm path length) was 0.44. C8 reverse phase HPLC and SDS PAGE both indicated a purity of approx 80%. The latter technique showed the major band had an apparent molecular weight of 23 000, clearly shifted from the original parent molecular weight of 21 000; on reduction the 23 000 band shifted to two bands with molecular weights of approx 21 000 and approx 5 000. The lyophilisate was easily soluble in 50mM formic acid or in PBS 'A' (Dulbecco) at a protein concentration of 6mg/ml.
- (d) [SCR1-3]-Cys-S-S-[MSWP-1] from (c) was divided into 0.3 ml aliquots and freeze-dried. Individual aliquots were resolubilised in 50mM formic acid (0.3ml or 0.039ml).

Example 9 Preparation of [SCR1-3/switch fusion]disulphide linked to [MAET]
(SEQ ID NO: 31)




- Title compound can be synthesised using TNB-activated SCR1-3/switch (SEQ ID NO: 7) prepared as in Example 7(d). The TNB-activated SCR1-3/switch is mixed with a molar excess of MAET (Example 1), which might be typically made up at 2.0mg/ml in DMSO, equivalent to about 3mM free thiol. Typical reaction conditions would be 1 to 4 hours at room temperature or overnight at 4 degrees C using a protein concentration of 1 to 100 uM. The reaction may be monitored by checking the generation of yellow colour, which is caused by the release of free TNB ion. Once the reaction is complete the solution may be buffer exchanged into a suitable buffer, for example 0.1M sodium phosphate pH 7.0, and stored at - 40 degrees C until required.

$$\text{H}_2\text{N} \cdot [\text{SCR 1-3}] \cdot \text{Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Pro-Ser-Lys-Ser-Ser-Gly-Cys-OH}$$


0.02ml of MSWP-1 (Example 2, 10mM in 0.1M sodium phosphate pH 7.0) was mixed with 0.005ml of TCEP (20mM in 50mM Hepes) and left for 10 min at room temperature. The resultant solution was Solution A containing the myristoylated peptide of SEQ ID NO: 5. TNB-activated SCR1-3/switch (SEQ ID NO: 7) prepared in a similar way to that described in Example 7(c) (0.3ml; 15uM in 0.1M sodium phosphate pH 7.0) was mixed with 0.0056ml of Solution A to give a theoretical MSWP-SH molar excess of five-fold over protein. The mixture was left for 4h at room temperature followed by 18h at 4 degrees C. Analysis by SDS PAGE followed by protein staining indicated one major band at apparent M_r 23K, corresponding to unreacted protein, and a minor band at apparent M_r 26K, corresponding to title protein.

25 **TNB-activated SCR1-3/switch product (SEQ ID NO: 7) (10uM; 0.43ml)**
prepared in a similar way to that described in Example 7(d) was mixed with TCEP
(5mM; 0.0026ml) and incubated for 17h at room temperature to yield the free fusion
protein SEQ ID NO: 7. MSWP-1 (10mM; 0.0086ml) was added and incubation was
continued for a further 4h. Small particles or crystals were present in the solution, but it
30 was otherwise clear. The particulate solution was buffer-exchanged into 50mM formic
acid (1.0ml), aliquoted and frozen. Analysis by SDS PAGE under non-reducing
conditions showed a number of bands, which included a species with an apparent
molecular weight of 25 000 - the target species.


Example 11 Preparation of [CR1: 1-1929]-Cys-S-S-[MSWP-1] (SEQ ID NO: 10)

10 Human complement receptor 1 (CR1, CD35) is a known regulator of complement activation which has been produced in a recombinant soluble form containing all of the extracellular SCR domains of a major natural allotype (Fearon et al, WO 89/09220, WO 91/05047). This form (sCR1) has been expressed as an active protein in Chinese Hamster Ovary (CHO) cells. Mutagenesis of the DNA sequence immediately downstream of the codon for Cys-1924 is performed to generate a new C-terminal cysteine residue.

15 A suitable example of a modified terminus of the cDNA sequence of sCR1 is as follows:

(5909) Bal I (5914)
 20CCT CTG GCC AAA TGT ACC TCT CGT GCA CAT TGC TGA
 The codon Asp-1930 in CR1 is replaced by that for a Cysteine (followed by a stop codon to generate a soluble protein) through ligation of a modified oligonucleotide to the unique Bal I restriction endonuclease site at position 5914 (numbering from Fearon et al, 1989,1991).

25 Expression of this modified cDNA in CHO cells and isolation of the product by standard chromatographic procedures generates a modified sCR1 protein which can be treated as in Example 8(a), (b) or (c) to couple it to MSWP-1 (Example 2) to yield the title compound.

Example 12 Preparation of [SCR1-3]-Cys-S-S-[MSWP-2] (SEQ ID No. 11)

40 [SCR1-3]-Cys protein (SEQ ID NO: 6) prepared in a similar way to that described in Example 6(d) (46uM protein; 0.20ml) was mixed with TCEP (5mM;


0.0054ml) and incubated at room temperature for approx. 20h. 0.05ml of this solution was mixed with 0.025ml of 0.1M ethanolamine and 0.003ml of MSWP-2 (see Example 3; 5mM in DMSO;); the solution was incubated for a further 3h at room temperature. SDS PAGE analysis showed the major band in the preparation had an apparent molecular weight of 23 000, clearly shifted from the original parent molecular weight of 21 000. The purity of the target protein was estimated from the SDS PAGE gel to be approx 80%.

Example 13 Preparation of [SCR1-3]-Cys-S-S-[MSWP-3] (SEQ ID No. 12)



[SCR1-3]-Cys protein (SEQ ID NO: 6) prepared in a similar way to that described in Example 6(d) (46uM protein; 0.10ml) was mixed with TCEP (5mM; 0.0037ml) and incubated at room temperature for approx. 18h. 0.01ml of 0.5M ethanolamine was added. 0.03ml of this 0.11ml solution was mixed with 0.0032ml of MSWP-3 (see Example 4; 2mM in 0.1M sodium phosphate pH 7.0); the solution was incubated for a further 3h at room temperature. SDS PAGE analysis showed the major band in the preparation had an apparent molecular weight of 23 000, clearly shifted from the original parent molecular weight of 21 000. The purity of the target protein was estimated from the SDS PAGE gel to be approx 80%.


Example 14 Preparation of [SCR1-3]-Cys-S-S-[TCPT-1] (SEQ ID No. 13)



[SCR1-3]-Cys protein prepared in a similar way to that described in Example 6(d) (46uM protein; 0.08ml) was mixed with TCEP (5mM; 0.0029ml) and incubated at room temperature for approx. 18h. 0.008ml of 0.5M ethanolamine was added. 0.04ml of this 0.088ml solution was mixed with 0.0029ml of TCPT-1 (see Example 5; 2.9mM in


DMSO). The TCPT-1 was added in 6 aliquots over a 2h period to minimise aggregation. The solution was incubated for a further 2h at room temperature. The final appearance of the mixture was one of a colloidal suspension and centrifugation at 2000g for 1 min showed that the target protein was compartmentalised in the precipitate. SDS PAGE analysis showed the major band in the preparation had an apparent molecular weight of about 23 000, clearly shifted from the original parent molecular weight of 21 000. The purity of the target protein was estimated from the SDS PAGE gel to be approx 80%.

Example 15 Preparation of a Rabbit anti-(human erythrocyte membrane) antibody - [MSWP-1] conjugate (RAEM-MSWP-1) (SEQ ID NO: 32)



S


S



Rabbit polyclonal anti(human erythrocyte membrane) (RAEM) antiserum (Dako, Denmark, 13mg/ml, 0.25ml) was diluted to 1.0ml with 50mM sodium phosphate 0.1M sodium chloride pH 7.4 (PBS) and treated with 30 ul of 100mM 2-iminothiolane in PBS (freshly dissolved) for 30 min at 25°C. These conditions have been shown (R.A.G.Smith & R.Cassels, Fibrinolysis, 2,189-195, 1988) to introduce an average of 2-3 free thiol groups per molecule of immunoglobulin G.


The product was purified by gel permeation chromatography on a small disposable column of Sephadex G-25m (PD-10, Pharmacia, Stockholm, Sweden) at 4°C. 2.5ml of the product (total volume 3.0ml, theoretical protein concentration ~6.1 uM) was treated with MSWP-1 (Example 2, 0.125 ml of 5mM solution in dimethyl sulfoxide, final conc ~240uM) and incubated at 25°C for 30 min. The product was gel-filtered on a PD10 column as above to give 3.0ml of a solution ~5uM in protein. This was stored frozen at -70°C.

Example 16 Preparation of a conjugate of Streptokinase and MSWP-1 (SEQ ID No 21)



S


S



Streptokinase (SK) stock solution (Behringwerke, Marburg, Germany, 12.8 mg/ml, 271 uM, 2.5ml) was gel filtered using a PD10 column into 3.2 ml of PBS buffer (see Example 15) containing 0.01%w/v Tween 80 [PST buffer]. Freshly made up 2-
 5 iminothiolane (64ul of 100mM) was added and the mixture incubated at 25°C for 1h. The product was gel filtered in 2 x 1.6ml batches into 2x 3.0 ml PST at 4°C on two PD10 columns. This solution was stored in aliquots of 1.5ml at -75°C.

Titration of the product with Ellman's reagent (0.1mM in 0.5ml 0.1M Triethanolamine.HCl pH 8.0) showed that it contained approximately 0.3mM free thiol
 10 groups. This corresponds to an average of 3- 3.5 thiol groups per molecule of SK. The stock thiolated SK solution (2 x 0.5ml) was processed by modifying one aliquot with MSWP-1 (32ul of 5mM stock in DMSO), incubated 1h at 25°C and gel filtered (PD10 column) into 3.0ml PST at 4°C. A control aliquot was processed in parallel without exposure to MSWP-1. Both products contained ~ 0.8 mg/ml protein based on an
 15 extinction coefficient of $0.76 \text{ (mg/ml)}^{-1}$ at 280nm for SK and were stored at -75°C.

Example 17 Reversible linkage of MSWP-1 to the active centre of Human Tissue-type Plasminogen Activator (SEQ ID No 22)



The thiol-reactive acyl-enzyme 4-N-[2-N-(3-[2-pyridyldithio]-
 25 ethylcarbonyl)aminoethyl]aminobenzoyl - [Ser-478] human tissue-type plasminogen activator [PDAEB->tPA] was prepared by the method of Smith and Cassels (Fibrinolysis, 2, 189-195, 1988). Tissue plasminogen activator (Actilyse, Boehringer
 Ingelheim, Germany, approx 2mg) was dissolved in the PST buffer of Example 16 (1.0ml) and treated with 25ul of a 20mM solution of 4'-amidinophenyl 4-N-[2-N-(3-[2-
 30 pyridyldithio]-ethylcarbonyl)aminoethyl]aminobenzoate hydrochloride (S.B.Kalindjian & R.A.G.Smith, Biochem. J. 248, 409-413, 1987) in dimethylsulphoxide. The mixture was incubated for 1h at 25°C and stored frozen at -80°C. It was reduced by addition of dithiothreitol (5ul of 0.5M in water) for 30 min at 0°C followed by buffer-exchange into PST buffer (3.0ml) as described in Example 16. The product was divided immediately
 35 into a retained sample (0.6ml) and a reaction sample (2.4ml) which was mixed with MSWP-1 (Example 2, 100ul of a 5mM solution in dimethylsulphoxide) and incubated for 90 min on ice. The product was buffer-exchanged as above into 3.2ml PST and stored in aliquots at -196°C.

Example 18 Expression and purification of [SCR1-3]-Cys (SEQ ID 6) from a fermentation run.

(a) Fermentation of *E. coli* harbouring the plasmid pDB1030

A frozen stock of *E. coli* harbouring the plasmid pDB1030 was initially prepared by plating the culture out onto LB agar plus ampicillin at 100µg/ml. 1ml aliquots were preserved in a 10% glycerol / PBS cryopreservative and stored under liquid nitrogen. A 1ml vial was thawed and was used to inoculate 100ml LB^{Amp¹⁰⁰} primary seed medium (Difco Bactotryptone, 10gl⁻¹; Difco yeast extract, 5gl⁻¹; sodium chloride, 5gl⁻¹; pH pre-sterilisation 7.4) in a 500ml flask. The primary seed stage was incubated at 37°C for 3 hours before transfer to the second seed stage, also 100ml LB^{Amp¹⁰⁰} per 500ml flask using a 1% inoculum. Following incubation as above for a further 4 hours a 1% inoculum was transferred to the tertiary seed stage, 10litres LB^{Amp¹⁰⁰} in a 15litre Biolafitte fermenter. The 10 litres tertiary seed medium was sterilised *in situ* for 45 minutes at 121°C before inoculation. Following incubation for 14.5 hours, the tertiary seed was transferred to the final stage fermenter as a 6% inoculum. Incubation conditions for the seed stage were as follows: airflow at 10lmin⁻¹ (1.0vvm), temperature 37°C, agitation at 400rpm (1.9ms⁻¹) and overpressure 0.2bar. 300 litres Tryptone phosphate medium^{Amp¹⁰⁰} (Difco Bactotryptone, 20gl⁻¹; Difco yeast extract, 15gl⁻¹; sodium chloride, 8gl⁻¹; disodium hydrogen orthophosphate, 2gl⁻¹; potassium dihydrogen orthophosphate, 1gl⁻¹; Dow Corning 1520 antifoam, 0.1gl⁻¹; pH pre-sterilisation 7.4) was sterilised *in situ* for 30 minutes at 121°C in a 450L Bioengineering fermenter. The fermenter was inoculated with 20 litre inoculum from the tertiary seed stage and incubated under the following conditions : airflow 450L min⁻¹ (1.5vvm), temperature 37°C, agitation 230rpm (1.5ms⁻¹) and overpressure 0.5bar. After an OD_{550nm} of 3.5 was obtained, 1mM IPTG was added. Harvest followed after continued incubation for 2 hours. A cell slurry was recovered after primary centrifugation through a Westfalia CSA19 (two discharges). The cells were further spun at 4700rpm (7000g) for 30 minutes in a Sorvall RC3B centrifuge. The total cell yield (wet weight) was 2.98Kg and was stored at -80 degrees C in approx. 600g lots.

(b) Isolation of inclusion bodies and purification of [SCR1-3]-Cys

Inclusion bodies from 100 g (wet weight) cell pellet were isolated and solubilised essentially as described in Example 6. The purification of target protein from resolubilised inclusion bodies was also as described in Example 6 with some modifications. The major ones were :

1. The use of Macroprep High S (Biorad) instead of S-Sepharose. 200g of matrix was used for 100g of cell pellet that had been sonicated. 1.4g of approx. 60%

pure target protein was produced in the solubilised and partially purified fraction on bodies.

2. Refolding of a 100mg sample of the partially purified protein was carried out by diluting the fully denatured protein (2mg/ml) 100-fold in cold 60mM ethanolamine/1mM EDTA, followed by addition of the glutathione redox couple.

The product of the above process was capable of being modified with MSWP-1 (Example 2) in a way similar to that described in Example 8.

Example 19 Expression and isolation of [SCR1-3(delN195-K196)]TNANKSLSSISCQT (SEQ ID NO: 14)

(a) Construction of plasmid pBC04-29 encoding [SCR1-3(delN195-K196)]TNANKSLSSISCQT

- Plasmid pBC04-29 was constructed from plasmid pDB1013-5 encoding SCR1-3 of LHR-A of CR1 (patent application WO 94/00571) by QuickChange site directed mutagenesis (Stratagene) according to the manufacturers protocols. Two complementary oligonucleotides (SEQ ID No 15 and SEQ ID No 16) were used to generate a novel restriction site (silent) at G186/P187 and a C terminal cysteine. In the event the reaction produced a frame-shift mutation at position N195. In the resulting sequence the C terminal amino acids N195 and K196 are replaced with a 14 amino acid peptide TNANKSLSSISCQT. Fortuitously, this sequence contains an internal cysteine close to the C terminus, preceded by a spacer sequence of 11 amino acids.

(b) Expression of plasmid pBC04-29 encoding [SCR1-3(delN195-K196)]TNANKSLSSISCQT in *E. coli*

- pBC04-29 was transformed into competent *E. coli* BL21(DE3)pLys-S and resultant colonies were isolated and checked for plasmid content. A single colony was inoculated into 10 ml LB medium (10g/L bactotryptone, 5g/L yeast extract, 10g/L NaCl) containing 50ug/ml ampicillin. The culture was grown for 6 - 18 hours at 37°C, 230 r.p.m. before being used to inoculate 1 litre of the same medium containing 50 ug/ml ampicillin at a dilution of 1 in 100 in 4 L erlenmeyer flasks. Cells were grown to an OD of 0.8 - 1.0 at A₆₀₀ nm. IPTG (isopropyl B-D galactopyranoside) was added to a final concentration of 1 mM and cells allowed to continue growth for a further 3 - 4 hours or overnight before harvesting by centrifugation at 8000 g/10 min. Pellet from 1L of culture was stored at -80°C.

(c) **Isolation and purification of [SCR1-3(delN195-K196)]TNANKSLSSISCQT**

The methods are essentially those detailed in Dodd I. et al (1995) Protein Expression and Purification 6 727-736, subsequently modified as described in Example 18. Briefly, the cell pellet from 1L of culture from (b) was resuspended in buffer, sonicated and the inclusion bodies isolated by centrifugation. The inclusion bodies were resolubilised in 100 ml of fully reducing buffer and target protein purified on Macroprep High S (30g wet weight). Product (27ml at nominal 1.5mg/ml) that eluted from the column in the 1M NaCl-containing buffer was refolded by dilution into 2.5L cold 60mM ethanolamine/1mM EDTA, with the glutathione redox agents added at 1h. After 3d at 4 degrees C the solution was ultrafiltered using a YM10 membrane and the retentate was treated with ammonium sulphate, centrifuged and the supernatant purified on Butyl Toyopearl 650M (bed volume 53ml). A single A280 peak was eluted by the decreasing ammonium sulphate gradient. SDS PAGE under non-reducing conditions followed by protein staining revealed a major polypeptide with an apparent molecular weight of 22 000, believed to be the target protein. One of the contaminating polypeptides had an apparent molecular weight of about 40 000, which was identified as the dimer of the monomeric form of the target by comparison with adjacent markers of [SCR1-3]-Cys. The product had an estimated protein concentration of 30uM.

20 **Example 20 Preparation of [SCR1-3(delN195-K196)]TNANKSLSSISC(-S-S-[MSWP-1])QT (SEQ ID No. 17)**

H₂N-[SCR 1-3(delN195-K196)]-Thr-Asn-Ala Asn- Lys-Ser-Leu-Ser-Ser-Ile-Ser-Cys-Gln-Thr

25 |
S
|
S
|

N-(myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Lys-Pro-Gly-Asp-Cys-NH₂

30 [SCR1-3(delN195-K196)]TNANKSLSSISCQT prepared as described in Example 19 (approx. 30uM protein; 0.1ml) was mixed with TCEP (5mM in 50mM Hepes pH 4.5; 0.0072ml) and incubated at room temperature (22 degrees C) for 15h. 0.05ml of this solution was mixed with 0.005ml of 0.5M ethanolamine and 0.003ml of 7mM MSWP-1 (see Example 2); the solution was incubated for a further 4h at room temperature. SDS
35 PAGE analysis showed a major band in the preparation had an apparent molecular weight of 25 000, clearly shifted from the original parent molecular weight of 23 000.

Example 21 Preparation of [SCR1-3]DGPSEILRGDFSSC (SEQ ID No. 23)**(a) Construction of plasmid pBC04-31 encoding [SCR1-3]DGPSEILRGDFSSC**

Plasmid pBC04-31 was constructed using plasmid pBC04-29 (described in Example 19) and a synthetic oligonucleotide pair (SEQ ID No. 25 and SEQ ID No. 26).
 5 pBC04-29 was digested with the restriction enzymes HindIII and ApaI and the large fragment (2170bp) isolated. The two oligonucleotides were annealed by warming to >90°C and slowly cooling to room temperature and ligated with the DNA fragment. The ligated DNA was transformed into competent *E. coli* XLI-Blue. Colonies were analysed for plasmids in which the oligonucleotides had been inserted by looking for the
 10 presence of a novel AvaI site at position 2733. On digestion with AvaI pBC04-31 yielded fragments of 2311 and 495bp. DNA from positive clones was used to transform the expression strains. The oligonucleotides inserted added the peptide sequence DGPSEILRGDFSSC to the C terminus of SCR1-3 and also repaired the frame-shift error seen in pBC04-29.

15

(b) Expression, isolation and purification of [SCR1-3]DGPSEILRGDFSSC

Expression, isolation and purification of [SCR1-3]DGPSEILRGDFSSC is carried out using pBC04-31 by procedures generally described in Example 6.

20

Example 22 Preparation of [SCR1-3]DGPSEILRGDFSSC-(S-S-[MSWP-1]) (SEQ ID No. 24)

H₂N-[SCR 1-3]-Asp-Gly-Pro-Ser-Glu-Ile-Leu-Arg-Gly Asp-Phe-Ser-Ser-Cys-OH

25

|
S
|
S
|

N-(myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Pro-Gly-Asp-Cys-NH₂

30

[SCR1-3] DGPSEILRGDFSSC protein prepared in a similar way to that described in Example 21 is reacted with MSWP-1 as described in Example 8 to give the title compound.

35 Example 23 Preparation of [SCR1-3] AAPSVIGFRILLKLVAGC (SEQ ID No. 33)**(a) Construction of plasmid pBC04-34 encoding [SCR1-3] AAPSVIGFRILLKLVAGC**

Plasmid pBC04-34 was constructed using plasmid pBC04-29 (described in Example 19) and a synthetic oligonucleotide pair (SEQ ID No. 34 and SEQ ID No. 35). pBC04-29 was digested with the restriction enzymes HindIII and ApaI and the large fragment (2170bp) isolated. The two oligonucleotides were annealed by warming to
 5 >90°C and slowly cooling to room temperature and were ligated with the DNA fragment. The ligated DNA was transformed into competent *E. coli* XLI-Blue. Colonies were analysed for plasmids in which the oligonucleotides had been inserted by looking for an increase in size of the NdeI/HindIII fragment by 59 base pairs. The presence of the
 10 cysteine codon was determined by the presence of a DdeI site at position 2781. pBC04-34 digested with DdeI yielded diagnostic bands of 481 and 109bp. DNA from positive clones was used to transform the expression strains (see next section). The
 oligonucleotides inserted added the peptide sequence AAPSVIGFRILLKLVAGC to the
 C terminus of SCR1-3 and also repaired the frame-shift error seen in pBC04-29.

- 15 (b) **Expression, isolation and purification of [SCR1-3]APSVIGFRILLKLVAGC**
 Expression isolation and purification of [SCR1-3] APSVIGFRILLKLVAGC is carried out using pBC04-34 by procedures generally described in Example 6.

Example 24 Preparation of [SCR1-3]AAPSVIGFRILLKLVAGC -(S-S-[MSWP-1])
 20 (SEQ ID No. 36)

H₂N-[SCR 1-3]-Ala-Ala-Pro-Ser-Val-Ile-Gly-Phe-Arg-Ile-Leu-Leu-Leu-Lys-Val-Arg-Gly-Cys-OH

|

S

|

S

|

25

N-(myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Lys-Pro-Gly-Asp-Cys-NH₂

[SCR1-3]AAPSVIGFRILLKLVAGC protein prepared in a similar way to that
 30 described in Example 23 is reacted with MSWP-1 as described in Example 8.

Biological Activity**(I) Anti-complement Activity Measured by the Classical Pathway-mediated Haemolysis of Sheep Erythrocytes**

(i) Functional activity of complement inhibitors was assessed by measuring the inhibition of complement-mediated lysis of sheep erythrocytes sensitised with rabbit antibodies (Diamedix Corporation, Miami, USA). The assay is designed to be specific for the classical pathway of complement activation. Human serum diluted 1:500 or 1:400 (final concentration in assay mixture) in 0.1 M Hepes/0.15 M NaCl/0.1% gelatin pH 7.4 was used as a source of complement. The serum was prepared from a pool of volunteers essentially as described in Dacie & Lewis, 1975. Briefly, blood was warmed to 37°C for 5 minutes, the clot removed and the remaining serum clarified by centrifugation. The serum fraction was split into small aliquots and stored at -196°C or -80°C. Aliquots were thawed as required and diluted in the Hepes buffer immediately before use.

Inhibition of complement-mediated lysis of sensitised sheep erythrocytes was measured using a standard haemolytic assay using a v-bottom microtitre plate format as follows:

50 µl of a range of concentrations of inhibitor diluted in Hepes buffer were mixed with 50 µl of the diluted serum and 100 µl of sensitised sheep erythrocytes and then incubated for 1 hour at 37°C. Samples were spun at 1600rpm at ambient temperature for 3 minutes before transferring 150 µl of supernatant to a flat bottom microtitre plate and determining the absorption at 405 or 410 nm. Maximum lysis (A_{max}) was determined by incubating serum with erythrocytes in the absence of any inhibitor. Background lysis (A_0) was determined by incubating erythrocytes in the absence of any serum or inhibitor. Inhibition was expressed as a fraction of the total cell lysis such that IC_{50} represents the concentration of inhibitor required to give 50% inhibition of lysis.

$$\% \text{ inhibition} = 1 - [(A - A_0) / (A_{max} - A_0)]$$

Results

Compound	IH50
WO94/00571 SEQ ID NO 27	0.2 - 0.3 ug/ml [10-15 nM] (1)
Example 6*	0.65 ug/ml [30nM] (mean of two) (2)
Example 7*	0.3-1.0 ug/ml [15-50nM] (n = 3)
Example 8a	0.014 ug/ml [0.6 nM])
Example 8b	<0.001 ug/ml [< 0.04nM]) (3)
Example 8c	0.001 ug/ml [0.043 nM])
Example 8d ⁺	[0.06nM])
Example 10a	0.02 ug/ml [0.8nM]
Example 10b	~0.01 ug/ml [~0.4nM]
Example 12	~0.0016 ug/ml [0.07 nM]
Example 13	~0.009 ug/ml [0.4nM]
Example 14	~1.1 ug/ml [50nM]
Example 19	[4nM]

* As 2-mercaptoethanol/glutathione derivatives

+ Assay of the two solutions and the original pre-lyophilisation solution from Example

5 8d.

Other IH₅₀ values generated for similar batches include:

- (1) 15nM
- (2) 8nM, 5nM, 8nM, 4nM
- (3) 0.3nM, 0.2nM, 0.07nM, 0.06nM, 0.2nM, 0.4nM, 0.5nM, 0.6nM.

10

The above data show that:

1. The complement inhibitory activities of the 'base' protein (SCR1-3 of human complement receptor 1 of WO94/00571) and its derivatives with either an additional C-terminal cysteine (SCR1-3/cys, Example 6) or a single cationic 'switch' sequence (SCR1-3/switch, Example 7) are similar.
- 15 2. However, incorporation of two membrane binding elements (electrostatic switch and myristoyl) by addition of MSWPs-1, 2 or 3 (which contain both elements) to SCR1-3/cys or three membrane binding elements by addition of the MSWP-1 to the SCR1-3/switch construct results in products which are significantly more potent (~20-200x)
- 20 than the base or single membrane binding element proteins. The use of TCTP-1 which is targeted to membrane elements found in CD3-positive cells and not to erythrocyte membranes gave a conjugate of similar potency to SCR1-3 derivatives with no or single

membrane addresses. Thus, the increases in potency in an assay which depends on an erythrocyte membrane event (cytolysis by the membrane attack complex of complement) can be attributed to membrane targeting of the cytolysis inhibitor proteins by the combination of two membrane binding elements.

5

(ii) Assay of anti-complement activity in the classical pathway haemolytic assay: activity in the sera of domestic pig, guinea pig, rat and marmoset.

The activity of [SCR1-3]-Cys-S-S-[MSWP-1] was examined in the classical pathway haemolytic assay using the sera of pig, guinea pig, rat or marmoset. The methodology was essentially as described in (I) with minor modifications, for example small changes to the concentration of serum used. [SCR1-3]-Cys-S-S-[MSWP-1] was prepared essentially as described in Example 8c. The IH50 values for the different sera were: pig, 0.2nM; guinea pig, 0.3nM; rat, 0.4nM; marmoset, 0.2nM. These results show that [SCR1-3]-Cys-S-S-[MSWP-1] is capable of inhibiting classical pathway complement inhibition in the sera of a variety of animal species.

10
15

(II) Anti-complement Activity Measured by Alternative Pathway-mediated Haemolysis of Guinea Pig Erythrocytes

Functional activity of complement inhibitors was assessed by measuring the inhibition of complement mediated lysis of guinea pig erythrocytes essentially as described by Scesney, S. M. et al (1996) J. Immunol. 26 1729-1735. The assay is designed to be specific for the alternative pathway of complement activation. Human serum prepared from a pool of volunteers essentially as described in Dacie & Lewis, 1975 was used as the source of complement. Briefly, blood was warmed to 37°C for 5 minutes, the clot removed and the remaining serum clarified by centrifugation. The serum fraction was split into small aliquots and stored at -196°C or -80°C. Aliquots were thawed as required and diluted in 0.1 M Hepes/ 0.15 M NaCl / 0.1% gelatin / 8 mM EGTA / 5 mM MgCl₂ pH 7.4 (buffer A) immediately before use. Guinea pig erythrocytes were prepared from guinea pig whole blood collected into EDTA-coated tubes as follows. The blood was spun at 1600 rpm for 5 min and the erythrocyte pellet washed 3 times with 0.1 M Hepes/ 0.15 M NaCl / 0.1% gelatin pH 7.4 until the supernatant of the spin was essentially colourless. The erythrocytes were finally resuspended to the original volume of blood used and were stored at + 4 degrees C. They were used within 2 weeks.

50 µl of a range of concentrations of inhibitor diluted in buffer A in a v-bottom microtitre plate were mixed with, first, 100 µl of serum that had been diluted 1:3 and second, 50 µl of guinea pig erythrocytes (diluted 1:49 in buffer A) and incubated for 1

20
25
30
35

hour at 37°C. The plate was spun at 1600 rpm for 3 minutes before transferring 150 µl of each supernatant to a flat bottom microtitre plate and determining the absorption at 405 nm, which reflects the amount of lysis in each test solution. Maximum lysis (A_{max}) was determined by incubating serum with erythrocytes in the absence of any inhibitor.

- 5 Background lysis (A_o) was determined by incubating erythrocytes in the absence of any serum or inhibitor. The final dilution of serum used in the assay did absorb at 405nm but the level of absorbance (approx 10% of A_{max}) was considered to have a negligible affect on the overall assay results and it was ignored in the calculations. Inhibition was expressed as a fraction of the total cell lysis such that IH50 represents the concentration
- 10 of inhibitor required to give 50% inhibition of lysis.

$$\% \text{ inhibition} = 1 - [(A - A_o) / (A_{\text{max}} - A_o)]$$

Results

- 15 Two aliquots (one lyophilised and resolubilised in a neutral buffer, the other not lyophilised) of a single batch of [SCR1-3]-Cys-S-S-[MSWP-1] prepared in a similar way to that described in Example 8 (c) were tested in the haemolytic assay. The IH50 values for the compounds were:

	[SCR1-3]-Cys-S-S-[MSWP-1] (not lyoph)	310nM
20	[SCR1-3]-Cys-S-S-[MSWP-1] (lyoph)	480nM

The result shows that [SCR1-3]-Cys-S-S-[MSWP-1] exhibited activity against the alternative pathway of the complement system and that lyophilisation and subsequent resolubilisation of the protein had no affect (within experimental error) on the biological activity of the protein.

25

(III) Plasminogen activator assay

- (i) SK-related molecules from Example 16 were assayed using a plasminogen activation assay. A solution of purified human Lys₇₇-Plasminogen (1µM in PST buffer containing 25%v/v glycerol [PSTG buffer], 0.5ml) was incubated with thiolated SK
- 30 (final concentration 0.1 to 1.07 nM) for 1h at 25°C. An aliquot of this mixture (10ul) was incubated with 1.0mM of the plasmin substrate S-2251 (H-D-Val-Leu-Lys-p-nitroanilide, KabiVitrum, Stockholm, Sweden) in 0.1M Triethanolamine HCl pH 8.0 (0.5ml) at 25°C. The release of p-nitroaniline was monitored continuously at 405nm. Under these conditions, one substrate unit (SU) of plasmin activity is defined as the
- 35 amount of enzyme giving an increase in optical density at 405nm of 0.001 min⁻¹. Under these conditions thiolated SK (1nM) generated plasmin at a nearly linear of 4225 SU/ml.

SK-MSWP-1 conjugate was diluted 1:100 in PSTG buffer and 5-50ul aliquots tested in the plasminogen activation assay. The stock preparation was found to contain approximately 2.9uM functional SK.

- (ii) The potential activity of the acyl-enzyme preparations of Example 17 was estimated by dilution 25-50 fold into PST buffer and incubation for 2h at 37°C, followed by assay using 2mM S-2288 (H-D-Ile-Pro-Arg-p-nitroanilide 2HCl) under the same conditions used in (i) above. Under these conditions, the potential activity of the reduced PDAEB->tPA preparation was 2760 SU/ml and that of the MSWP-1/PDAEB->tPA conjugate 535 SU/ml.

10

(IV) Erythrocyte binding assays

(i) Erythrocyte aggregation test for modified and unmodified Rabbit anti-(human erythrocyte membrane) antibody.

- Human pooled erythrocytes (Ortho A2, Raritan, New Jersey, 3% v/v, 50ul) were added to microtitre plates wells and either unmodified rabbit anti-(human erythrocyte membrane) antibody [RAEM] or RAEM-MSWP1 conjugate from Example 15 added at concentrations expressed relative to undiluted stock RAEM. Cells were agitated at ~100rpm for 40min at 25°C. 5ul was removed from each well and examined by light microscopy at x 20 magnification. A visual scoring scale was used as follows:

20

- No clumping, cells moving freely relative to each other.
- + Small clumps (<10 cells)
- ++ Larger clumps (100 plus cells)
- +++ Very large visible aggregates

25 Results

Controls (n=6)	-	RAEM-MSWP1 1/3900	+/-
RAEM 1/1100	-	RAEM-MSWP1 1/1000	+/-
RAEM 1/600	-		
RAEM 1/350	+/-	RAEM-MSWP1 1/357	+++
RAEM 1/50	++	RAEM-MSWP1 1/62	+++

Conclusion

- The antibody preparation modified to contain a membrane-binding unit was more effective at inducing aggregation of cells because binding to the cell membrane through MSWP1 allowed a higher effective concentration of bridging antibody on the membrane surface.

30

(ii) Binding of 125-Iodine-[SCR1-3]-Cys-S-S-[MSWP-1] to human erythrocytes

[SCR1-3]-Cys-S-S-[MSWP-1] (2mg/ml in PBS; 0.25ml) was mixed with 0.5mCi of 125-Iodine (Amersham) in the presence of 9nmol potassium iodide following the Iodogen procedure and reagents (Pierce and Warriner (UK) Ltd.). The labelling was allowed to proceed for 20 min at room temperature, the reaction was quenched with 0.1ml of 1M potassium iodide and the solution buffer-exchanged into PBS/0.1% albumin. Citrated blood collected from a healthy volunteer was used as a source of human erythrocytes. Blood (0.2ml) was mixed with 10 microlitres of appropriately diluted 125-Iodine-[SCR1-3]-Cys-S-S-[MSWP-1] (final concentration 700pM) and incubated for 30min at 37 degrees C. The erythrocytes were then isolated by three repeat washings in PBS / centrifugation steps and samples counted in a Wallac 1470 Wizard gamma counter. The results were as follows:

	cpm
1st wash	3 600 000
1st pellet	140 000
2nd wash	52 000
3rd wash	6 500
final pellet	26 000

Using values of 5×10^9 erythrocytes per ml of blood and a specific radioactivity of 2.7×10^7 cpm / nmole for the [SCR1-3]-Cys-S-S-[MSWP-1] it was calculated that about 600 molecules of [SCR1-3]-Cys-S-S-[MSWP-1] bound per cell (the value for 'final pellet').

(iii) Binding of fluorescein-labelled-[SCR1-3]-Cys-S-S-[MSWP-1] to human erythrocytes

[SCR1-3]-cys (prepared in a similar way to that described in Example 18) (45uM, 1.0mg/ml in 0.1M sodium phosphate, approx. 0.2M ammonium sulphate pH 7.0) was partially reduced by incubation at 25°C for 18h by the addition of a 4-molar excess of Tris(2-carboxyethyl)phosphine (TCEP; Pierce & Warriner (UK) Ltd.). The solution was buffer exchanged into 50mM Hepes pH7.0; post buffer exchange the protein concentration was 22uM. The partially reduced [SCR1-3]-cys was incubated with a 4-fold molar excess of 6-(fluorescein-5-carboxamido)hexanoic acid, succinimidyl ester (Molecular Probes Inc., USA) and incubated for 1h at 4°C. The excess fluorescent label was removed by buffer exchange of the protein solution into 50mM Hepes pH7.0. Fluorescein-[SCR1-3]-cys-S-S-[MSWP-1] was synthesised by adding MSWP-1 (Example 2) to give a five-fold molar excess over fluorescein labelled protein and was

incubated for 4h at 25°C. The solution was buffered exchanged into PBS and this solution was used for the microscopy studies.

[SCR1-3], 10mg/ml in 50mM formic acid, was mixed in a 1:10 ratio with 50mM NaHCO₃ pH8.5; the pH of the solution was adjusted with NaOH to pH9.5. The fluorescein was extracted from Celite-fluorescein isothiocyanate (Celite:fluorescein; 1:10, Sigma) by DMSO in a 1:4 (w/v) ratio. The fluorescein-DMSO solution was added to the protein solution in a 1:14 ratio and incubated for 1h at RT. Excess label was removed by gel filtration into PBS containing 0.01% Tween-80 and this solution was used for microscopy studies.

Citrated blood was collected from a healthy volunteer and the erythrocytes isolated, washed in PBS and diluted 250-fold compared to the original blood volume. 0.05ml of erythrocytes were incubated with 2uM fluorescein-[SCR1-3]-cys-S-S-[MSWP-1] or 2uM fluorescein-[SCR1-3] and incubated for 30min at 37°C. An eight microlitre sample of each incubation was mounted on a slide and viewed on an inverted confocal microscope (Biorad). The cells incubated with fluorescein-[SCR1-3] showed no specific staining whereas with those incubated with fluorescein-[SCR1-3]-cys-S-S-[MSWP-1] staining appeared diffusely over the cell surface and also intensely stained patches were visible on the cell membrane. No labelling was seen intracellularly.

(iv) Binding of MSWP-1/PDAEB->tPA to human erythrocytes

Human trypsinized and glutaraldehyde-treated red blood cells (1.0ml of a 4% suspension) was pelleted by low-speed centrifugation and resuspended in a total volume of 0.5ml PST containing either no additions or approximately 270 SU of either reduced PDEAB->tPA or MSWP-1/ PDAEB->tPA conjugate of Example 17. The mixtures were incubated by gentle rolling for 5 min at 23°C and then the cells were pelleted by centrifugation followed by two washes with 1.0ml PST buffer. Finally, the cells were suspended in 0.5ml PST and incubated at 37°C. Samples of the supernatant (100ul) were removed after pelleting. Assay using S-2288 (as above) showed that after 2h, approximately 7% of the applied t-PA activity was present in the supernatant of cells exposed to MSWP-1/PDAEB->tPA whereas only ~2.8% was present in the supernatant of cells exposed to reduced PDAEB->tPA alone. No t-PA amidolytic activity was detected in controls.

This experiment suggests that reversible linkage of the active site of t-PA to MSWP-1 increases the tendency of this enzyme to bind to red blood cells.

(v) **Localisation of SK-MSWP-1 conjugate on the surface of human erythrocytes**

A stabilised preparation of human erythrocytes (trypsinised, glutaraldehyde-treated, Sigma, Gillingham, UK, 4% v/v, 0.4ml) was pelleted by centrifugation (~2000g/2min) and resuspended in 0.4ml PST buffer with either 0.1uM thiolated SK or 0.1uM SK-MSWP-1 from Example 16.

The suspensions were incubated for 30 min at 37°C and then washed by two cycles of centrifugation and resuspension in PST buffer. Finally, they were resuspended in PSTG buffer (0.4ml) containing 1uM plasminogen and incubated and assayed for plasmin as described above.

The control thiolated-SK generated plasmin at a rate of 522 SU/ml, while the SK-MSWP-1 conjugate produced 6184 SU/ml. The latter activity corresponds to around 2100 thiolated SK molecules/cell.

(vi) **Binding of [SCR1-3]-Cys-S-S-[MSWP-1] to human erythrocyte membranes**

4 X 2.0ml of trypsinized, glutaraldehyde-treated human erythrocytes (Sigma, R0127) were centrifuged for 2min at about 3000 rpm. The supernatants were discarded and the cells resuspended in phosphate/saline/Tween (0.01%) (PST) (1ml per tube) and [SCR1-3]-Cys-S-S-[MSWP-1] of Example 8 was added to a final concentration of 20ug/ml to three of the tubes. The mixtures were then incubated at 37 degrees C for 30 min., then washed five times by repeat centrifugation and washing in PST. The cells were finally suspended in 1ml PST and were held at 4 degrees C.

The ability of these cells to inhibit complement-mediated lysis of sheep erythrocytes was measured using the standard classical pathway complement inhibition assay described in (I) above. The human erythrocytes were added to the assay at four different dilutions, followed by the human serum and then the sheep red blood cells and incubation at 37 degrees C as usual. The % inhibition data are shown below.

Final dilution	human erythrocytes, untreated	human erythrocytes, treated with [SCR1-3]-Cys-S-S-[MSWP1]
1/4	22%	62%
1/16	-8%	88%
1/256	5%	74%
1/2500	-7%	51%

Thus the percentage inhibition for the [SCR1-3]-Cys-S-S-[MSWP-1]-treated cells at maximum dilution was significantly higher than the untreated cells at 1/4 dilution. The [SCR1-3]-Cys-S-S-[MSWP-1]-treated cells, therefore, contained at least 600-fold more complement inhibitory activity than the untreated cells, even though the cells had been washed extensively to remove any non-bound [SCR1-3]-Cys-S-S-[MSWP-1].

TABLE

SEQUENCE LISTING

(<- next to a peptide sequence in {} signifies sequence runs C to N terminus)

5

SEQ ID NO:1:

GCACCGCAGTGCATCATCCCGAACAAATGCTAATAAA

SEQ ID NO:2:

10 AGCTTTTATTAGCATTTGTTCTGGGATGATGCACTGCG

SEQ ID NO:3:

GCACCGCAGTGCATCATCCCGAACAAAGACGGTCCGAAAAAGAAGAAAAAGAAATCTCCGTCCAAATCTTCC
GGTTGCTAATAAA

15

SEQ ID NO:4:

AGCTTTTATTAGCAACCGGAAGATTTGGACGGAGATTTCTTTTCTTCTTTTCCGACCGTCTTTGTTCTGGG
ATGATGCACTGCG

20

SEQ ID NO:5:

Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Lys-Pro-Gly-Asp-Cys-NH₂

SEQ ID NO:6:

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
25 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
30 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
35 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
Ile Ile Pro Asn Lys Cys

SEQ ID NO:7:

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 5 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 10 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 Ile Ile Pro Asn Lys Asp Gly Pro Lys Lys Lys Lys Lys Lys Ser Pro
 15 Ser Lys Ser Ser Gly Cys

SEQ ID NO:8:

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 20 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 25 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 30 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 -> <-
 Ile Ile Pro Asn Lys Cys-S-S-(Cys Asp Gly Pro Lys Lys Lys Lys Lys
 | |
 CO₂H CONH₂
 35 Lys Ser Pro Ser Lys Ser Ser Gly) (N-Myristoyl)

SEQ ID NO:9:

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 5 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 10 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 Ile Ile Pro Asn Lys Asp Gly Pro Lys Lys Lys Lys Lys Lys Ser Pro
 -> <-
 15 Ser Lys Ser Ser Gly Cys-S-S-(Cys Asp Gly Pro Lys Lys Lys Lys Lys
 | |
 CO₂H CONH₂
 Lys Ser Pro Ser Lys Ser Ser Gly) (N-Myristoyl)

20 SEQ ID NO 10
 linear, 2 polypeptide chains disulphide linked
 1 [QCNAPWLPF ARPTNLTDEF EFPIGTYLNY ECRPGYSGRP
 41 FSIICLKNSV WTGAKDRCRR KSCRNPPDPV NGMVHVIKGI
 81 QFGSQIKYSC TKGYRLIGSS SATCIISGDT VIWDNETPIC
 25 121 DRIPCGLPPT ITNGDFISTN RENFHYGSVV TYRCNPGSGG
 161 RKVFELVGEP SIYCTSNDQ VGIWSPAPQ CIIPNKCTPP
 201 NVENGILVSD NRSFLSLNEV VEFRCQPGFV MKGPRRVKQ
 241 ALNKWEPELP SCSRVCQPPP DVLHAERTQR DKDNFSPGQE
 281 VFYSCEPGYD LRGAASMRCT PQGDWSPAAP TCEVKSDDF
 30 321 MGQLLNGRVL FPNLQLGAK VDFVCDEGFQ LKGSSASYCV
 361 LAGMESLWNS SVPVCEQIFC PSPVIPNGR HTGKPLEVFP
 401 FGKAVNYTCD PHPDRGTSFD LIGESTIRCT SDPQNGVWS
 441 SPAPRCGILG HCQAPDHFLF AKLKTQTNAS DFPIGTSILKY
 481 ECRPEYYGRP FSITCLDNLV WSSPKDVCKR KSKTPPDPV
 35 521 NGMVHVITDI QVGSRINYSC TTGHRLLIGHS SAECILSGNA
 561 AHWSTKPPIC QRIPCGLPPT IANGDFISTN RENFHYGSVV
 601 TYRCNPGSGG RKVFELVGEP SIYCTSNDQ VGIWSPAPQ
 641 CIIPNKCTPP NVENGILVSD NRSFLSLNEV VEFRCQPGFV

681 MKGPRRVKCQ ALNKWEPELP SCSRVCQPPP DVLHAERTQR
 721 DKDNFSPGQE VFYSCEPGYD LRGAASMRCT PQGDWSPAAP
 761 TCEVKSDDF MGQLLNGRVL FPNLQLGAK VDFVCEGFQ
 801 LKGSSASYCV LAGMESLWNS SVPVCEQIFC PSPPVIPNGR
 5 841 HTGKPLEVFP FGKAVNYTCD PHPDRGTSFD LIGESTIRCT
 881 SDPQNGVWS SPAPRCGILG HCQAPDHFLF AKLKTQTNAS
 921 DFPIGTSILKY ECRPEYYGRP FSITCLDNLV WSSPKDVCKR
 961 KSCKTPDPDV NGMVHVITDI QVGSRIYNSC TTGHRLLIGHS
 1001 SAECILSGNT AHWSTKPPIC QRIPCGLPPT IANGDFISTN
 10 1041 RENFHYGSVV TYRCNLGSRG RKFVFLVGEF SIYCTSNDDQ
 1081 VGIWSGPAPQ CIIPNKCTPP NVENGILVSD NRSFLSLNEV
 1121 VEFRCQPGFV MKGPRRVKCQ ALNKWEPELP SCSRVCQPPP
 1161 EILHGEHTPS HQDNFSPGQE VFYSCEPGYD LRGAASLHCT
 1201 PQGDWSPEAP RCAVKSCDDF LGQLPHGRVL FPLNLQLGAK
 15 1241 VSFVCEGFR LKGSSVSHCV LVGMRLWNN SVPVCEHIFC
 1281 PNPPAILNGR HTGTPSGDIP YGKEISYTCF PHPDRGMTFN
 1321 LIGESTIRCT SDPHGNGVWS SPAPRCESV RAGHCKTPEQ
 1361 FPFASPTIPI NDFEFPVGTG LNYECRPGYF GKMFSISCLF
 1401 NLVWSSVEDN CRRKSCGPPP EPFNGMVHIN TDTQFGSTVN
 20 1441 YSCNEGFRLI GSPSTTCLVS GNNVTWDKKA PICEIISCEP
 1481 PPTISNGDFY SNNRTSFHNG TVVTYQCHTG PDGEQLFELV
 1521 GERSIYCTSK DDQGVWSSP PPRCISTNKC TAPEVENAIR
 1561 VPGNRSFFSL TEIIRFCQP GFVMVGSHTV QCQTNGRWGP
 1601 KLPHCSRVCQ PPPEILHGEH TLSHQDNFSP GQEVFYSECF
 25 1641 SYDLRGAASL HCTPQGDWSP EAPRCTVKSC DDFLGQLPHG
 1681 RVLLPLNLQL GAKVSFVCFE GFRLKGRSAS HCVLAGMKAL
 1721 WNSSVPVCEQ IFCPNPPAIL NGRHTGTPFG DIPYKGEISY
 1761 ACDTHPDRGM TFNLIGESSI RCTSDPQNG VWSSPAPRCE
 1801 LSVPAACPHP PKIQNGHYIG GHVSLYLPGM TISYTCDFGY
 30 1841 LLVGKGFIFC TDQGIWSQLD HYCKEVNCSF PLFMNGISKE
 1881 LEMKKVYHYG DYVTLKCEDG YTLEGSPWSQ CQADDRWDPP

 -> <-
 1921 LAKCTSRAC]-S-S-(CDGPKKKKKKSPSKSSG)-(N-Myristoyl)
 | |
 35 CO₂H CONH₂

In SEQ ID NO 10, peptide sequences are given in brackets in single letter amino acid code.

SEQ ID No. 11

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 5 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 10 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 15 Ile Ile Pro Asn Lys Cys-S-S-Cys-Asp-Gly-Pro-Lys-Lys-Lys-Lys-

$$\begin{array}{c} | \\ \text{NHCOCH}_3 \end{array}$$
Lys-Ser-Pro-Ser Lys-Ser-Ser-Lys-(N-Myristoyl)-NH₂

20 SEQ ID No. 12

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 25 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 30 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys

->

<-

35 Ile Ile Pro Asn Lys Cys-S-S-(Cys-Lys-Thr-Lys-Ser-Lys-Lys-Lys-Lys-

$$\begin{array}{c} | \\ \text{CONH}_2 \end{array}$$

Lys-Lys-Gly-Asp-Lys-Ser)-NH-(Myristoyl)

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
5 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
10 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
15 Ile Ile Pro Asn Lys Cys-S-S-Cys-Ser-Ala-Ala-Pro-Ser-Ser-Gly-Phe-
NHCOCH₃
Arg-Ile-Leu-Leu-Leu-Lys-Val-NH(CH₂)₉CH₃

20 SEQ ID NO:14

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
25 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
30 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
Ile Ile Pro Thr Asn Ala Asn Lys Ser Leu Ser Ser Ile Ser Cys Gln
Thr

35

SEQ ID No 15

CTGGAGCGGGCCCGCACCGCAGTGCATCATCCCGAACAAATGCTAATAAAAGC

SEQ ID No 16

GCTTTTATTAGCATTTGTTTCGGGATGATGCACTGCGGTGCGGGCCCGCTCCAG

SEQ ID No 17

5 linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 10 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 15 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys

20 Ile Ile Pro Thr Asn Ala Asn Lys Ser Leu Ser Ser Ile Ser Cys-S-S-(Cys-
 | |
 Gln-Thr CONH₂
 -Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Lys-Ser-Ser-Gly)-NH-(Myristoyl)

SEQ ID No 18

25 Cys-Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Pro-Ser-Lys-Ser-Ser-Lys-NH₂

SEQ ID No 19

Ser-Lys-Asp-Gly-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Lys-Thr-Lys-Cys

30 SEQ ID No 20

Cys-Ser-Ala-Ala-Pro-Ser-Ser-Gly-Phe-Arg-Ile-Leu-Leu-Leu-Lys-Val

SEQ ID No 21

linear, 2 polypeptide chains disulphide linked

35 N-(myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Pro-
 Gly-Asp-Cys-NH₂

|
 S-S-[4-butyrimino]-N-ε(Lys) [Streptokinase]

SEQ ID No 22

Single chain form of the 527 amino acid residue intact t-PA molecule.

Residue 478 (serine) has been modified as shown below

- [SYQVICRDEKTMIIYQQHQSWLRPVLRSNRVEYCWNSGRAQCHSVPVKSCSEPRCFN
 5 GGTCQQALYFSDFVCQCEGAFAGKCCCEIDTRATCYEDQGISYRGTWSTAESGAECTNW
 NSSALAQKPYSGRRPDAIRLGLGNHNYCRNPDRDSKPWCYVFKAGKYSSEFCSTPACS
 EGNSDCYFGNGSAYRGTHSLTESGASCLPWNSMILIGKVYTAQNPSAALGLGKHNYC
 RNPDGDAKPWCHVLKNRRLTWEYCDVPSCSTCGLRQYSQPQFRIKGLFADIASHPWQA
 AIFAKHRRSPGERFLCGGILISSCWILSAAHCFQERFPPHLLTVILGRTYRVVPGEE
 10 EQKFEVEKYIVHKEFDDDTYDNDIALQLKSDSSRCAQESSVVRTVCLPPADLQLPDW
 TECELSGYGKHEALSPFYSERLKEAHVRLYPSSRCTSQHLLNRTVTDNMLCAGDTRSG
 GPQANLHDACQGDSCGGLVCLNDGRMTLVGIIISWGLGCGQKDVPGVYTKVTNYLDWIRDNMRI
 15 Ser 478 O-4-CO-benzyl-NH(CH₂)₂NHCO(CH₂)₂-S-S-(Cys-Asp-Gly-Pro-Lys-Lys-
 |
 CONH₂
 Lys-Lys-Lys-Lys-Ser-Pro-Ser-Lys-Ser-Ser-Gly)-NH-(Myristoyl)

SEQ ID No 23

- 20 Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 25 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 30 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 Ile Ile Pro Asn Lys Asp Gly Pro Ser Glu Ile Leu Arg Gly Asp Phe
 Ser Ser Cys

35 SEQ ID No 24

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys

Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 5 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 10 Ile Ile Pro Asn Lys Asp-Gly-Pro-Ser-Glu-Ile-Leu-Arg-Gly Asp-Phe-

<-

Ser-Ser-Cys-S-S-(Cys-Asp-Gly-Pro-Lys-Lys-Lys-Lys Lys-Lys-Ser-Pro-Ser-
 Lys-Ser-Ser-Gly)-NH-(Myristoyl)

15 SEQ ID No 25
 CGCACCGCACTGCATCATCCCGAACAAAGATGGCCCGAGCGAAATTCTGCGTGGCGATTTTAGCAGCTGCTA

SEQ ID No 26
 ACGTTAGCAGCTGCTAAAATCGCCACGCAGAATTTGCTCGGGCCATCTTTGTTCTGGGATGATGCACTGCGG
 20 TGCGGGCC

SEQ ID No 27
 N-(Myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Lys-Pro-
 Gly-Asp-(S-2-thiopyridyl)Cys-NH₂

25 SEQ ID No 28
 N-acetyl-(S-2-thiopyridyl)Cys Asp-Gly-Pro-Lys-Lys-Lys-Lys-Lys-Lys-Ser-
 Pro-Ser Lys-Ser-Ser-(εN-(Myristoyl))Lys-NH₂

30 SEQ ID No 29
 N-(Myristoyl)-Ser-Lys-Asp-Gly-Lys-Lys-Lys-Lys-Lys-Lys-Ser-Lys-Thr-Lys-
 (S-2-Thiopyridyl)Cys-NH₂

SEQ ID No 30
 35 N-acetyl-(S-2-thiopyridyl)Cys-Ser-Ala-Ala-Pro-Ser-Ser-Gly-Phe-Arg-Ile-
 Leu-Leu-Leu-Lys-Val-NH(CH₂)₉CH₃

SEQ ID No 31

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 5 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 10 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 Ile Ile Pro Asn Lys Asp Gly Pro Lys Lys Lys Lys Lys Lys Ser Pro
 15 Ser Lys Ser Ser Gly Cys-S-S-(CH₂)₂-CONH-(CH₂)₁₂CH₃

SEQ ID No 32

linear, 2 polypeptide chains disulphide linked

N-(myristoyl)-Gly-Ser-Ser-Lys-Ser-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Lys-Pro-

20 Gly-Asp-Cys-NH₂

S-S-[4-butyrimino]-N-ε(Lys){Rabbit anti-(human erythrocyte
 membrane) antibody}

25 SEQ ID No 33

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
 Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
 Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
 30 Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
 Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
 Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
 Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
 35 Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
 Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
 Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
 Ile Ile Pro Asn Lys Ala Ala Pro Ser Val Ile Gly Phe Arg Ile Leu
 Leu Leu Lys Val Gly Cys

SEQ ID No 34

CGCACCGCAGTGCATCATCCCGAACAAGCGGCGCCAGCGTGATTGGCTTCCGTATTCTGCTGCTGAAAGT
GGCGGGCTGCTA

5

SEQ ID No 35

AGCTTAGCAGCCCGCCACTTTCAGCAGCAGAATACGGAAGCCAATCACGCTGGGCGCCGCTTTGTTCGGGAT
GATGCACTGCGGTGCGGGCC

10 SEQ ID No 36

linear, 2 polypeptide chains disulphide linked

Met Gln Cys Asn Ala Pro Glu Trp Leu Pro Phe Ala Arg Pro Thr Asn
Leu Thr Asp Glu Phe Glu Phe Pro Ile Gly Thr Tyr Leu Asn Tyr Glu
Cys Arg Pro Gly Tyr Ser Gly Arg Pro Phe Ser Ile Ile Cys Leu Lys
15 Asn Ser Val Trp Thr Gly Ala Lys Asp Arg Cys Arg Arg Lys Ser Cys
Arg Asn Pro Pro Asp Pro Val Asn Gly Met Val His Val Ile Lys Gly
Ile Gln Phe Gly Ser Gln Ile Lys Tyr Ser Cys Thr Lys Gly Tyr Arg
Leu Ile Gly Ser Ser Ser Ala Thr Cys Ile Ile Ser Gly Asp Thr Val
Ile Trp Asp Asn Glu Thr Pro Ile Cys Asp Arg Ile Pro Cys Gly Leu
20 Pro Pro Thr Ile Thr Asn Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn
Phe His Tyr Gly Ser Val Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly
Gly Arg Lys Val Phe Glu Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr
Ser Asn Asp Asp Gln Val Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys
Ile Ile Pro Asn Lys Ala Ala Pro Ser Val Ile Gly Phe Arg Ile Leu

25

<-

Leu Leu Lys Val Gly Cys S-S-(Cys-Asp-Gly-Pro-Lys-Lys-Lys-Lys Lys-Lys-
Ser-Pro-Ser-Lys-Ser-Ser-Gly)-NH-(Myristoyl)

CLAIMS

1. A soluble derivative of a soluble polypeptide, said derivative comprising two or more heterologous membrane binding elements with low membrane affinity covalently associated with the polypeptide which elements are capable of interacting, independently and with thermodynamic additivity, with components of cellular or artificial membranes exposed to extracellular fluids.
2. A derivative according to claim 1 wherein each membrane binding element with low membrane affinity has a dissociation constant $1\mu\text{M}$ - 1mM .
3. A derivative according to claim 1 or 2 wherein each membrane binding element has a size $<5\text{kDa}$.
4. A derivative according to any preceding claim which incorporates sufficient elements with low affinities for membrane components to result in a $0.01 - 10\text{nM}$ dissociation constant affinity for specific membranes.
5. A derivative according to any preceding claim which has a solubility in pharmaceutical formulation media $>100\mu\text{g/ml}$.
6. A derivative according to any preceding claim wherein at least one element is hydrophilic.
7. A derivative according to any preceding claim which comprises two to eight membrane binding elements.
8. A derivative according to any preceding claim wherein the membrane binding elements are selected from: fatty acid derivatives; basic amino acid sequences; ligands of known integral membrane proteins; sequences derived from the complementarity-determining region of monoclonal antibodies raised against epitopes of membrane proteins; and membrane binding sequences identified through screening of random chemical or peptide libraries.
9. A derivative according to claim 8 wherein a membrane binding element is a fatty acid derivative selected from aliphatic acyl groups with about 8 to 18 methylene units, long-chain (8-18 methylene) aliphatic amines and thiols, steroid and farnesyl derivatives.

10. A derivative according to claim 8 or 9 wherein a membrane binding element is a basic aminoacid sequence including (Lys)_n where n is from 3 to 10.

11. A derivative according to claim 10 wherein the amino acid sequence is selected from:

i) DGPKKKKKKSPSKSSG

ii) GSSKSPSKKKKKKPGD

iii) SPSNETPKKKKKRFSFKKSG

iv) DGPKKKKKKSPSKSSK

10 v) SKDGKKKKKKSKTK

(N-terminus on left)

12. A derivative according to any of claims 8 to 10 wherein a membrane binding element is a ligand of a known integral membrane protein selected from GRGDSP, 15 DGPSEILRGDFSS, GNEQSFRVLDLRTLRYA, GFRILLKKV, SAAPSSGFRILLKKV and AAPSVIGFRILLKKVAG or the carbohydrate ligand Sialyl Lewis^x.

13. A derivative according to any preceding claim wherein the soluble polypeptide is selected from IL-4 Y124D mutein, prourokinase, streptokinase, tissue-type plasminogen 20 activator, reteplase, leptin, complement inhibitors selected from complement regulatory proteins and hybrids or muteins thereof, scFv antibody against cytokines, Protein C, antibodies against CD4, B7/CD28, CD3/TCR or CD11b(CR3) and Interferon-β and derivatives.

25 14. A derivative according to any preceding claim which has the following structure:



in which:

P is the soluble polypeptide,

each L is independently a flexible linker group,

30 each W is independently a peptidic membrane binding element,

n is an integer of 1 or more and

X is a peptidic or non-peptidic membrane-binding entity which may be covalently linked to any W.

35 15. A derivative according to claim 14 wherein peptidic membrane binding elements are 8 to 20 amino acids long and elements W are located sequentially either at the N or C terminus of the soluble polypeptide and the amino acid sequences are linked to one

another and to the soluble peptide by linker groups which are selected from: hydrophilic and/or flexible aminoacid sequences of 4 to 20 aminoacids; linear hydrophilic synthetic polymers; and chemical bridging groups.

- 5 16. A derivative according to claim 14 or 15 wherein the chemical bridging groups are of formula (I):



- 10 in which each of A and B, which may be the same or different, represents -CO-, -C(=NH₂⁺)-, maleimido, -S- or a bond and R is a bond or a linking group containing one or more -(CH₂)- or meta-, ortho- or para- disubstituted phenyl units optionally together with a hydrophilic portion.

- 15 17. A derivative according to claim 16 wherein R is selected from -(CH₂)_r-, -(CH₂)_p-S-S-(CH₂)_q- and -(CH₂)_p-CH(OH)-CH(OH)-(CH₂)_q-, in which r is an integer of at least 2; and p and q are independently integers of at least 2, or (CH₂)₂CONH(CH₂)_nNH-(4-phenyl) where n is an integer of 3 to 8.

18. A soluble derivative according to claim 1 of SEQ ID NO: 32.

- 20 19. A soluble derivative according to any of claims 1 to 17 wherein the soluble polypeptide is a soluble complement inhibitor.

20. A soluble derivative according to claim 19 wherein the soluble polypeptide is a soluble CR1 polypeptide fragment.

- 25 21. A soluble derivative according to claim 20 wherein the soluble CR1 polypeptide consists of residues 1-196 of CR1 and with an N-terminal methionine and the derivative comprises a myristoyl group and one or more polypeptides sequence selected from

DGPKKKKKKSPSKSSGC

30 GSSKSPSKKKKKKPGDC

CDGPKKKKKKSPSKSSK

SKDGKKKKKSKTKC

CSAAPSSGFRILLKLV

AAPSVIGFRILLKVAGC

- 35 and

DGPSEILRGDFSSC

(N-terminus on left).

22. A soluble derivative according to claim 20 or 21 selected from SEQ ID NOs: 8, 31, 9, 10, 11, 12, 13, 17, 24 and 36.
- 5 23. A soluble derivative according to any of claims 1 to 17 wherein the soluble polypeptide is a thrombolytic agent.
24. A soluble derivative according to claim 23 selected from SEQ ID NOs: 21 and 22.
- 10 25. A process for preparing a derivative according to claim 1 which process comprises expressing DNA encoding the polypeptide portion of said derivative in a recombinant host cell and recovering the product and thereafter post translationally modifying the polypeptide to chemically introduce membrane binding elements.
- 15 26. A polypeptide portion of a derivative according to claim 1, comprising the soluble peptide linked by a peptide bond to one peptidic membrane binding element
27. A soluble polypeptide including a C-terminal cysteine.
- 20 28. The polypeptide of SEQ ID NO: 7, 23, 33, 6 or 14.
29. A DNA polymer encoding the polypeptide portion of claim 26, 27 or 28.
- 25 30. A replicable expression vector capable, in a host cell, of expressing the DNA polymer of claim 29.
31. A host cell transformed with a replicable expression vector of claim 30.
- 30 32. A peptide membrane binding element with low membrane affinity comprising one or more derivatisations selected from:
a terminal cysteine residue optionally activated at the thiol group;
an N-haloacetyl group (where halo signifies chlorine, bromine or iodine) located at the N-terminus of the the peptide or at an ϵ -amino group of a lysine residue;
35 an amide group at the C-terminus;
an N-terminal blocking group; and
a fatty acid N-acyl group at the N-terminus or at an ϵ -amino group of a lysine residue.

33. A peptidic membrane binding element derivatised according to claim 32 wherein the peptide has the amino sequence of a peptide defined in claim 11 or 12 and a fatty acid N-acyl group of 8 to 18 methylene units at the N-terminus or at an ϵ -amino group of a lysine residue of the peptide.
34. A peptidic membrane binding element derivatised according to claim 32 or 33 selected from SEQ ID Nos: 27, 28, 29 and 30.
35. A C₁₀₋₂₀ fatty acyl derivative of an aminoC₂₋₆alkane thiol (optionally C-substituted).
36. A compound according to claim 35 selected from N-(2-myristoyl) aminoethanethiol and N-myristoyl L-cysteine.
37. A pharmaceutical composition comprising a derivative according to any of claims 1 to 24 in combination with a pharmaceutically acceptable carrier.
38. A derivative according to any of claims 1 to 24 for use as a medicament.
39. A method of treatment of disorders amenable to treatment by a soluble peptide which comprises administering a soluble derivative of said soluble peptide according to any of claims 1 to 24.
40. The use of a derivative according to any of claims 1 to 24 for the preparation of a medicament for treatment of disorders amenable to treatment by the soluble peptide.
41. A pharmaceutical composition for treating a disease or disorder associated with inflammation or inappropriate complement activation comprising a therapeutically effective amount of a derivative according to claim 19 and a pharmaceutically acceptable carrier or excipient.
42. A method of treating a disease or disorder associated with inflammation or inappropriate complement activation comprising administering to a subject in need of such treatment a therapeutically effective amount of a derivative according to claim 19.

43. The use of a derivative of claim 19 in the manufacture of a medicament for the treatment of a disease or disorder associated with inflammation or inappropriate complement activation.
- 5 44. A pharmaceutical composition for treating a disease or disorder associated with inflammation or inappropriate complement activation comprising a therapeutically effective amount of a soluble CR1 polypeptide derivative according to any of claims 20 to 22, and a pharmaceutically acceptable carrier or excipient.
- 10 45. A method of treating a disease or disorder associated with inflammation or inappropriate complement activation comprising administering to a subject in need of such treatment a therapeutically effective amount of a soluble CR1 polypeptide derivative of any one of claims 20 to 22.
- 15 46. The use of a soluble CR1 polypeptide derivative of any one of claims 20 to 22 in the manufacture of a medicament for the treatment of a disease or disorder associated with inflammation or inappropriate complement activation.
- 20 47. A pharmaceutical composition for treating thrombotic disorders comprising a therapeutically effective amount of a derivative according to claim 23 or 24 and a pharmaceutically acceptable carrier or excipient.
- 25 48. A method of treating thrombotic disorders comprising administering to a subject in need of such treatment a therapeutically effective amount of a derivative according to claim 23 or 24.
49. The use of a derivative according to claim 23 or 24 in the manufacture of a medicament for the treatment of thrombotic disorders.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, C07K 14/705, 16/28, 14/315, C12N 9/72, A61K 47/48, G01N 33/68, C07C 323/41, 323/59		A3	(11) International Publication Number: WO 98/02454 (43) International Publication Date: 22 January 1998 (22.01.98)
(21) International Application Number: PCT/EP97/03715 (22) International Filing Date: 8 July 1997 (08.07.97) (30) Priority Data: 9614871.3 15 July 1996 (15.07.96) GB (71) Applicant (for all designated States except US): ADPROTECH PLC [GB/GB]; Unit 3, Number 2 Orchard Road, Royston, Herts. SG8 5HD (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): SMITH, Richard, Anthony, Godwin [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). DODD, Ian [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). MOSSAKOWSKA, Danuta, Ewa, Irena [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: DAVIES, Jonathan, Mark; Reddie & Grose, 16 Theobalds Road, London WC1X 8PL (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 2 April 1998 (02.04.98)	
(54) Title: CONJUGATES OF SOLUBLE PEPTIDIC COMPOUNDS WITH MEMBRANE-BINDING AGENTS			
(57) Abstract Soluble derivatives of soluble polypeptides incorporating membrane binding elements, their use in therapy and methods and intermediates including peptide membrane binding elements. Preferred membrane-binding elements are fatty acid derivatives or polylysine-containing sequences.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/03715

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/11 C07K14/705 C07K16/28 C07K14/315 C12N9/72
A61K47/48 G01N33/68 C07C323/41 C07C323/59

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K G01N C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 114 787 A (CIBA-GEIGY) 1 August 1984 see page 49, line 13 - line 16 ---	35,36
X	DATABASE WPI Week 25 Derwent Publications Ltd., London, GB; AN 89-238911 XP002053042 "Modified animal hair powder heavy metal absorbents etc. is obtained by reacting mercaptan(s) or mercapto-modified silicone oil with side chains of cystine in hair" & JP 01 174 528 A (SEIWA KASEI CO LTD.) , 11 July 1989 see abstract --- -/-	35,36



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

23 January 1998

Date of mailing of the international search report

17.02.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Masturzo, P

INTERNATIONAL SEARCH REPORT

Internu al Application No
PCT/EP 97/03715

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	T BACON-BAGULEY ET AL.: "Thrombospondin binding to specific sequences within the A-alpha and B-beta chains of fibrinogen" JOURNAL OF BIOLOGICAL CHEMISTRY., vol. 265, no. 4, 5 February 1990, MD US, pages 2317-2323, XP002053040 cited in the application see page 2320, line 1 - line 10 ---	1-24
A	C G YEH ET AL.: "Recombinant soluble human complement receptor type I inhibits inflammation in the reversed passive Arthus reaction in rats" JOURNAL OF IMMUNOLOGY, vol. 146, no. 1, 1 January 1991, BALTIMORE US, pages 250-256, XP002053041 cited in the application see the whole document ---	1-34, 40-49
A	WO 94 06826 A (UNITED BIOMEDICAL INC.) 31 March 1994 see the whole document ---	1-24
A	WO 94 00571 A (SMITH KLINE BEECHAM) 6 January 1994 cited in the application see the whole document ---	1-34, 40-49
A	WO 93 22343 A (THE ROCKEFELLER UNIVERSITY) 11 November 1993 see the whole document -----	1-34, 37-49

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 97/03715

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/EP 97/03715

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

Claims Nos.: 1-20,22-35,37-49

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

The vague nature of the subject matter of the above claims does not permit a complete search for economical reasons. A complete search has therefore been performed only on the subject matter of claims 21 and 36; the search concerning the other claims was limited to the real examples provided in the text of the present application.

Remark : Although claims 39, 42, 45, 48 are directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/03715

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 114787 A	01-08-84	AU 569865 B	25-02-88
		AU 2374584 A	26-07-84
		CA 1247089 A	20-12-88
		DE 3485094 A	31-10-91
		DK 31684 A	26-07-84
		JP 6008316 B	02-02-94
		JP 59139348 A	10-08-84
		US 4666886 A	19-05-87
WO 9406826 A	31-03-94	US 5582968 A	10-12-96
		AU 5127693 A	12-04-94
		CA 2144435 A	31-03-94
		EP 0662082 A	12-07-95
		FI 951198 A	15-03-95
		JP 8500122 T	09-01-96
		NO 950977 A	14-03-95
		US 5639594 A	17-06-97
WO 9400571 A	06-01-94	EP 0649468 A	26-04-95
		JP 7508516 T	21-09-95
WO 9322343 A	11-11-93	US 5580563 A	03-12-96

Inis Page Blank (uspto)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

inis Page Blank (uspro)